

Parameters for which the Lawrence-Krammer representation is reducible

Claire Levailant and David Wales
cl@caltech.edu, dbw@caltech.edu

Caltech, MC 253-37, Pasadena, CA 91125

January 24, 2009

Abstract

We show that the representation, introduced by Lawrence and Krammer to show the linearity of Braid groups, is generically irreducible, but for that for some values of its two parameters when these are specialized to complex numbers, it becomes reducible. To do so, we construct a representation of degree $\frac{n(n-1)}{2}$ of the BMW algebra of type A_{n-1} inside the Lawrence-Krammer space. As a representation of the Braid group on n strands, it is equivalent to the Lawrence-Krammer representation where the two parameters of the BMW algebra are related to the two parameters of the Lawrence-Krammer representation. We give the values of the parameters for which the representation is reducible and give the proper invariant subspaces in some cases. We use this representation to show that for these special values of the parameters and other values, the BMW algebra of type A_{n-1} is not semisimple.

1 Introduction

1.1 Introduction and main results

In [5], Daan Krammer constructed a representation of the Braid group in order to show that it is linear. Since this representation was earlier introduced by Ruth Lawrence in [6], it is called the Lawrence-Krammer representation. In this paper, we examine a representation of degree $\frac{n(n-1)}{2}$ of the BMW algebra of type A_{n-1} in the Lawrence-Krammer space. As a representation of the Braid group on n strands, it is equivalent to the Lawrence-Krammer representation (abbreviated L-K representation). By studying this representation we show that the L-K representation is generically irreducible but that for some values of its two parameters when these are specialized to complex numbers, it becomes reducible. Throughout the paper, we let l , m and r be three nonzero complex

parameters, where m and r are related by $m = \frac{1}{r} - r$. We define $\mathcal{H}_{F,r^2}(n)$ as the Iwahori-Hecke algebra of the symmetric group $Sym(n)$ over the field $F = \mathbb{Q}(l, r)$ with generators $g_1, \dots, g_{n-1}$, that satisfy the Braid relations and the relation $g_i^2 + m g_i = 1$ for all i . Our definition is the same as the definition of [8] after the generators have been rescaled by a factor $\frac{1}{r}$. Our main result is as follows.

Theorem 1. (Main theorem)

Let n be an integer with $n \geq 3$ and let m, l and r be three nonzero complex parameters, where m and r are related by $m = \frac{1}{r} - r$. Assume that $\mathcal{H}_{F,r^2}(n)$ is semisimple, and so assume that $r^{2k} \neq 1$ for every integer $k \in \{1, \dots, n\}$.

When $n \geq 4$, the Lawrence-Krammer representation of the BMW algebra of type A_{n-1} with parameters l and m over the field $\mathbb{Q}(l, r)$ is irreducible, except when $l \in \{r, -r^3, \frac{1}{r^{2n-3}}, \frac{1}{r^{n-3}}, -\frac{1}{r^{n-3}}\}$, when it is reducible.

When $n = 3$, the Lawrence-Krammer representation of the BMW algebra of type A_2 with parameters l and m over the field $\mathbb{Q}(l, r)$ is irreducible, except when $l \in \{-r^3, \frac{1}{r^3}, 1, -1\}$, when it is reducible.

A consequence of this result and of the method that we use is the following.

Theorem 2.

Let n be an integer with $n \geq 4$ and let l, m and r be three nonzero complex parameters, where m and r are related by $m = \frac{1}{r} - r$.

Suppose $n \geq 4$. If $r^{2k} = 1$ for some $k \in \{2, \dots, n\}$ or if l belongs to the set of values $\{r, -r^3, \frac{1}{r^{n-3}}, -\frac{1}{r^{n-3}}, \frac{1}{r^{2n-3}}, -r^{2n-3}, r^{n-3}, -r^{n-3}, \frac{1}{r^3}, -\frac{1}{r^3}\}$, the BMW algebra of type A_{n-1} with parameters l and m over the field $\mathbb{Q}(l, r)$ is not semisimple.

(Case $n = 3$). If $r^4 = 1$ or $r^6 = 1$ or if $l \in \{-r^3, \frac{1}{r^3}, 1, -1\}$, the BMW algebra of type A_2 with parameters l and m over the field $\mathbb{Q}(l, r)$ is not semisimple.

In [11], Wenzl states that the BMW algebra of type A_{n-1} is semisimple except possibly if r is a root of unity or $l = r^n$, for some $n \in \mathbb{Z}$. Here, Theorem 2 gives instances of when the algebra is not semisimple. The result of this theorem is also contained in the recent work of Hebing Rui and Mei Si (see [10]). They use the representation theory of cellular algebras.

1.2 The method

We show that the action on a proper invariant subspace of the Lawrence-Krammer space must be an Iwahori-Hecke algebra action.

First, we study the Iwahori-Hecke algebra representations of small degrees and investigate whether they may occur inside the L-K space and if so for which values of l and r . We will denote by $\mathcal{V}^{(n)}$ the L-K space. We show that if there exists a one-dimensional invariant subspace inside $\mathcal{V}^{(n)}$, it forces the value $\frac{1}{r^{2n-3}}$ for l , except when $n = 3$ when it forces $l \in \{-r^3, \frac{1}{r^3}\}$. Conversely, for these values of l and r , there exists a one-dimensional invariant subspace of $\mathcal{V}^{(n)}$ and the representation is thus reducible. Similarly, we show that if there exists an irreducible $(n-1)$ -dimensional invariant subspace inside $\mathcal{V}^{(n)}$, it forces $l = \frac{1}{r^{n-3}}$ or $l = -\frac{1}{r^{n-3}}$ in the case when $n \neq 4$ and $l \in \{-r^3, \frac{1}{r^3}, -\frac{1}{r^3}\}$ in the case when $n = 4$. Conversely, for each of these values of l and r , there exists an irreducible $(n-1)$ -dimensional subspace of $\mathcal{V}^{(n)}$, which shows the reducibility of the representation in these cases as well.

Second, we identify a proper invariant subspace of $\mathcal{V}^{(n)}$ which is nontrivial when $l = r$ (case $n \geq 4$) or $l = -r^3$. This shows that the representation is also reducible in these cases.

Third, we study in detail the small cases $n \in \{3, 4, 5, 6\}$.

At last, when $n \geq 7$, we use a result from representation theory: the irreducible representations of $\mathcal{H}_{F,r^2}(n)$ have degrees $1, n-1, \frac{n(n-3)}{2}, \frac{(n-1)(n-2)}{2}$ or degree greater than $\frac{(n-1)(n-2)}{2}$, except in the case $n = 8$, when they have degrees $1, 7, 14, 20, 21$ or degrees greater than 21. We use this fact, and proceed by induction on $n \geq 5$ to show that if $\mathcal{V}^{(n)}$ is reducible, it forces $l \in \{r, -r^3, \frac{1}{r^{2n-3}}, \frac{1}{r^{n-3}}, -\frac{1}{r^{n-3}}\}$. To do so, we use the fact that if the dimension of a proper invariant subspace $\mathcal{W}$ of $\mathcal{V}^{(n)}$ is large enough, then the intersections $\mathcal{W} \cap \mathcal{V}^{(n-1)}$ and $\mathcal{W} \cap \mathcal{V}^{(n-2)}$ are nontrivial.

1.3 Definitions

1.3.1 The BMW algebra

We recall below the defining relations of the BMW algebra $B(A_{n-1})$ (or simply B) of type A_{n-1} with nonzero complex parameters l and m over the field $\mathbb{Q}(l, r)$, where r is a root of the quadratic $X^2 - mX + 1$. This algebra has two sets of $(n-1)$ elements, namely the invertible g_i 's that satisfy the Braid relations (1) and (2) and generate the algebra and the e_i 's that generate an ideal. For nodes i and j with $1 \leq i, j \leq n-1$, we will write $i \sim j$ if $|i-j| = 1$ and $i \not\sim j$ if $|i-j| > 2$.

$$g_i g_j = g_j g_i \quad \text{if } i \not\sim j \quad (1)$$

$$g_i g_j g_i = g_j g_i g_j \quad \text{if } i \sim j \quad (2)$$

$$e_i = \frac{l}{m} * (g_i^2 + m g_i - 1) \quad \text{for all } i \quad (3)$$

$$g_i e_i = l^{-1} e_i \quad \text{for all } i \quad (4)$$

$$e_i g_j e_i = l e_i \quad \text{if } i \sim j \quad (5)$$

We will also use some direct consequences of these defining relations (see [2], Proposition 2.1):

$$e_i g_i = l^{-1} e_i \quad \text{for all } i \quad (6)$$

$$g_i^2 = 1 - m g_i + m l^{-1} e_i \quad \text{for all } i \quad (7)$$

$$g_i^{-1} = g_i + m - m e_i \quad \text{for all } i \quad (8)$$

as well as the following "mixed Braid relations" (see [2], Proposition 2.3):

$$g_i g_j e_i = e_j e_i \quad \text{if } i \sim j \quad (9)$$

$$g_i e_j e_i = g_j e_i + m(e_i - e_j e_i) \quad \text{if } i \sim j \quad (10)$$

This algebra was shown by Morton and Wassermann to be isomorphic to the tangle algebra of Morton and Traczyk (see [9]). All the algebraic relations given in this paper have a geometric formulation in terms of tangles. In particular, we will use the tangles in §3.4.

1.3.2 The Lawrence-Krammer space

We now recall some terminology associated with root systems of type A_{n-1} . Let $M = (m_{ij})_{1 \leq i \leq j \leq n-1}$ be the Coxeter matrix of type A_{n-1} . Let $(\alpha_1, \dots, \alpha_{n-1})$ be the canonical basis of $\mathbb{R}^{n-1}$ and let's define a bilinear form B_M over $\mathbb{R}^{n-1}$ by:

$$B_M(\alpha_i, \alpha_j) = -\cos\left(\frac{\pi}{m_{ij}}\right)$$

By the theory in [1], B_M is an inner product that we will simply denote by $(|)$. Let r_i denote the reflection with respect to the hyperplane $\text{Ker}(\alpha_i|.)$ of $\mathbb{R}^{n-1}$ and so:

$$\forall x \in \mathbb{R}^{n-1}, r_i(x) = x - 2(\alpha_i|x) \alpha_i$$

Finally, let ϕ^+ denote the set of $\frac{n(n-1)}{2}$ positive roots:

$$\begin{aligned} \phi^+ = \{ & \alpha_1, \alpha_2, \alpha_2 + \alpha_1, \alpha_3, \alpha_3 + \alpha_2, \alpha_3 + \alpha_2 + \alpha_1, \dots, \\ & \alpha_{n-1}, \alpha_{n-1} + \alpha_{n-2}, \alpha_{n-1} + \alpha_{n-2} + \dots + \alpha_1 \} \end{aligned}$$

We define $\mathcal{V}^{(n)}$ as the vector space over $\mathbb{Q}(l, r)$ with basis the x_β 's, indexed by the positive roots $\beta \in \phi^+$. Thus, $\dim_F \mathcal{V}^{(n)} = |\phi^+| = \frac{n(n-1)}{2}$. The so-defined space $\mathcal{V}^{(n)}$ is the Lawrence-Krammer space.

To each positive root, we associate an element of the BMW algebra in the following way:

- To α_1 we associate e_1 .
- To $\alpha_i = r_{i-1} \dots r_1 r_i \dots r_2(\alpha_1)$, we associate the algebra element $g_{i-1} \dots g_1 g_i \dots g_2 e_1$, which after using the defining rules (1) and (9) above simplifies to $e_i \dots e_1$.
- To $\alpha_j + \dots + \alpha_i = r_j \dots r_{i+1}(\alpha_i)$ where $j \geq i+1$, we associate the algebra element $g_j \dots g_{i+1} e_i \dots e_1$.

2 The representation

2.1 The BMW left module

In what follows, F still denotes the field $\mathbb{Q}(l, r)$ and H denotes the Hecke algebra of the symmetric group $\text{Sym}(n-2)$ over the field F with generators $g_3, \dots, g_{n-1}$ and relations the Braid relations and the relations $g_i^2 + m g_i = 1$ for each i . As $r^2 + m r - 1 = 0$, our base field F is a one-dimensional H -module for the action given by $g_i.1 = r$ for every integer i with $3 \leq i \leq n-1$. We define B_1 as the quotient of two left ideals of B :

$$B_1 = B e_1 / < B e_i e_1 >_{i=3 \dots n-1}$$

Since e_i commutes with e_j for any $i \not\sim j$, we have for each node i with $3 \leq i \leq n-1$:

$$e_1(g_i^2 + m g_i - 1) = \frac{l}{m} e_1 e_i = 0 \text{ in } B_1$$

Then, B_1 is a right H -module. Thus, B_1 is a left B -module and a right H -module. Since F is an H -module, we get a left B -module by considering the tensor product

$$B_1 \otimes_H F$$

This B -module is precisely the left representation of B that we study in this paper. Its degree is $\frac{n(n-1)}{2}$ since by the forthcoming computations, we have:

$$B_1 = \text{Span}_F(e_1, e_2e_1, g_2e_1, e_3e_2e_1, g_3e_2e_1, g_3g_2e_1, \dots, \\ e_{n-1} \dots e_1, g_{n-1}e_{n-2} \dots e_1, g_{n-1} \dots g_2e_1).H$$

We will denote by $\mathcal{B}$ the spanning set above. The action of the g_i 's on the elementary tensors $b \otimes_H 1$, where $b \in \mathcal{B}$, was computed in section 2.2 below. These computations show in particular that if $G_1(n)$ denotes the matrix of the left action of g_1 on the vectors

$$e_1 \otimes_H 1, e_2e_1 \otimes_H 1, g_2e_1 \otimes_H 1, e_3e_2e_1 \otimes_H 1, g_3e_2e_1 \otimes_H 1, g_3g_2e_1 \otimes_H 1, \dots, \\ e_{n-1} \dots e_1 \otimes_H 1, g_{n-1}e_{n-2} \dots e_1 \otimes_H 1, g_{n-1} \dots g_2e_1 \otimes_H 1$$

of $B_1 \otimes_H F$, and if $\det G_1(n)$ denotes its determinant, we have:

$$\begin{aligned} \det G_1(3) &= -\frac{1}{l} \\ \det G_1(n) &= -r^{n-3} \det G_1(n-1) \quad \forall n \geq 4 \end{aligned}$$

Thus, the determinant of $G_1(n)$ is nonzero, which shows that these vectors are linearly independent.

We notice that there is a bijection between $\mathcal{B}$ and the set of positive roots ϕ^+ , as described in the previous section. Let's name this bijection u .

2.2 The action by the g_k 's

We describe further the representation by computing the action of the g_k 's on the elementary tensors $b \otimes_H 1$, where b is an algebra element in the spanning set $\mathcal{B}$. An element b of $\mathcal{B}$ is of the form:

$$g_j \dots g_{i+1}e_i \dots e_1 \quad \text{with } j > i \geq 1 \quad (\text{or simply } g_{j,i+1}e_{i,1}) \quad (I)$$

that we will refer to as of type (I), or of the form:

$$e_i \dots e_1 \quad \text{with } i \geq 1 \quad (\text{or simply } e_{i,1}) \quad (II)$$

referred to as of type (II).

For $i \geq j$, we set $g_{i,j} = g_i \dots g_j$ and $e_{i,j} = e_i \dots e_j$, where $g_{i,i}$ and $e_{i,i}$ are simply g_i and e_i respectively. When $i < j$, we define $g_{i,j}$ to be the identity.

In what follows, we fix i and j as in (I) and (II). There are several cases.

2.2.1 Action by g_{i-1} (Case A)

Let's first compute the action of g_{i-1} for $i \geq 2$ on elements of both types. We have for the type (I):

$$\begin{aligned} g_{i-1}.b &= g_{j,i+1}g_{i-1}e_{i,1} && \text{by (1)} \\ &= g_{j,i}e_{i-1,1} + m e_{i-1,1}g_{j,i+1} - m g_{j,i+1}e_{i,1} && \text{by (10) and (1)} \end{aligned}$$

And for the type (II):

$$g_{i-1}.b = g_i e_{i-1,1} + m e_{i-1,1} - m e_{i,1} \quad \text{by (10)}$$

Thus, we get:

$$g_{i-1}.(b \otimes_H 1) = \begin{cases} g_{j,i} e_{i-1,1} \otimes_H 1 + m r^{j-i} e_{i-1,1} \otimes_H 1 - m g_{j,i+1} e_{i,1} \otimes_H 1 \\ g_i e_{i-1,1} \otimes_H 1 + m e_{i-1,1} \otimes_H 1 - m e_{i,1} \otimes_H 1 \end{cases}$$

where the first line refers to type (I) and the second line to type (II). For future references, we name these two equalities $A(I)$ and $A(II)$ respectively.

2.2.2 Action by g_i (Case B)

We have for types (I) and (II) respectively:

$$g_i.b = g_{j,i+2} e_{i+1,1} \quad \text{by (9)}$$

$$g_i.b = l^{-1} e_{i,1} \quad \text{by (4)}$$

Thus, we get:

$$g_i.(b \otimes_H 1) = \begin{cases} g_{j,i+2} e_{i+1,1} \otimes_H 1 & B(I) \\ l^{-1} e_{i,1} \otimes_H 1 & B(II) \end{cases}$$

Notice that if $j = i + 1$, expression $B(I)$ reduces to $e_{i+1,1} \otimes_H 1$.

2.2.3 Action by g_j (Case C)

Let's first deal with Type (I). We have by (7):

$$g_j.b = g_{j-1,i+1} e_{i,1} - m g_{j,i+1} e_{i,1} + m l^{-1} e_j g_{j-1,i+1} e_{i,1}$$

We will rearrange the last term of the sum and to do so, we will need more mixed braid relations, as in the following lemma.

Lemma 1.

$$e_j e_i g_j = e_j g_i^{-1} \quad \text{when } i \sim j \quad (11)$$

$$e_i e_j e_i = e_i \quad \text{when } i \sim j \quad (12)$$

Proof. These are equalities (8) and (10) of Proposition 2.3 in [2]. $\square$

Using the relations of Lemma 1, we now give a new expression for $e_j g_{j-1,i+1} e_{i,1}$.

Lemma 2.

$$e_j g_{j-1,i+1} e_{i,1} = e_{j,1} g_j^{-1} \dots g_{i+2}^{-1} \quad (13)$$

Proof. Using Lemma 1, we replace e_j with $e_j e_{j-1} e_j$, then replace $e_{j-1} e_j g_{j-1}$ with $e_{j-1} g_j^{-1}$ to get:

$$e_j g_{j-1,i+1} e_{i,1} = e_j e_{j-1} g_{j-2,i+1} e_{i,1} g_j^{-1}$$

Proceeding inductively, we obtain (13). $\square$

When b is of the second type, we simply have:

$$g_j.b = \begin{cases} e_{i,1}g_j & \text{if } j > i + 1 \\ g_{i+1}e_{i,1} & \text{if } j = i + 1 \end{cases}$$

In the following expressions, the first line is for type (I) and the next two are for type (II).

$$g_j.(b \otimes_H 1) = \begin{cases} g_{j-1,i+1}e_{i,1} \otimes_H 1 - m g_{j,i+1}e_{i,1} \otimes_H 1 + \frac{m}{lr^{j-i-1}} e_{j,1} \otimes_H 1 \\ r e_{i,1} \otimes_H 1 & \text{when } j > i + 1 \\ g_{i+1}e_{i,1} \otimes_H 1 & \text{when } j = i + 1 \end{cases}$$

We will refer to these equations as (CI), (CII) and (CII') respectively.

2.2.4 Action by g_{j+1} (Case D)

Since

$$\begin{aligned} g_{j+1}.b &= g_{j+1}g_{j,i+1}e_{i,1} && \text{when } b \text{ is of type (I)} \\ g_{j+1}.b &= e_{i,1}g_{j+1} && \text{when } b \text{ is of type (II),} \end{aligned}$$

we simply get:

$$g_{j+1}.b = \begin{cases} g_{j+1,i+1}e_{i,1} \otimes_H 1 & (DI) \\ r e_{i,1} \otimes_H 1 & (DII) \end{cases}$$

2.2.5 Action by g_k where $k \notin \{i-1, i, j, j+1\}$ (Case E)

- Suppose first $k < i-1$ and b of type (I). We compute:

$$g_k.b = g_{j,i+1}e_{i,k+2}g_k e_{k+1}e_{k,1}$$

Expanding $g_k e_{k+1}e_k$ with (10) yields:

$$g_k.b = g_{j,i+1}e_{i,k+2}g_{k+1}e_{k,1} + m g_{j,i+1}e_{i,k+2}e_{k,1} - m g_{j,i+1}e_{i,1}$$

Since $e_{k+2}e_k = 0$ in B_1 , this expression simplifies as follows:

$$g_k.b = g_{j,i+1}e_{i,k+2}(g_{k+1} - m e_{k+1})e_{k,1}$$

Replacing $g_{k+1} - m e_{k+1} = g_{k+1}^{-1} - m$ with (8) and simplifying with $e_{k+2}e_k = 0$, we then obtain:

$$g_k.b = g_{j,i+1}e_{i,k+2}g_{k+1}^{-1}e_{k,1}$$

Applying equality (11) to $e_{k+2}g_{k+1}^{-1}$ now yields the new expression for $g_k.b$:

$$g_k.b = g_{j,i+1}e_{i,k+1}g_{k+2}e_{k,1},$$

which is also after commutation of g_{k+2} :

$$g_k.b = g_{j,i+1}e_{i,1}g_{k+2}$$

Thus, the action of g_k on the tensor $g_{j,i+1}e_{i,1} \otimes_H 1$ is simply a multiplication by r . After inspection, the computation for type (II) is identical and the action of g_k on the tensor $e_{i,1} \otimes_H 1$ is also a multiplication by r .

- Suppose now that $k > j + 1$. Visibly, g_k commutes with $g_{j,i+1}e_{i,1}$ and with $e_{i,1}$, so that in both cases, the action by g_k on the tensor $b \otimes_H 1$ is simply a multiplication by r .

- Finally, suppose k belongs to $\{i + 1, \dots, j - 1\}$ where $j \geq i + 2$. We look at the action of g_k on $g_{j,i+1}e_{i,1}$. We move g_k to the right, then use the Braid relation $g_k g_{k+1} g_k = g_{k+1} g_k g_{k+1}$, then move g_{k+1} this time to the end of the expression. After doing these moves, we get:

$$g_k g_{j,i+1} e_{i,1} = g_{j,i+1} e_{i,1} g_{k+1}$$

It follows in particular that

$$g_k \cdot b \otimes_H 1 = r b \otimes_H 1,$$

as in the previous cases. It remains to look at the action of g_k on $e_{i,1}$. We have:

$$g_k \cdot b = \begin{cases} g_{i+1} e_{i,1} & \text{if } k = i + 1 \\ e_{i,1} g_k & \text{otherwise} \end{cases}$$

We summarize Case E in the following two equalities:

$$\forall k \notin \{i-1, i, j, j+1\}, g_k \cdot b \otimes_H 1 = \begin{cases} g_{i+1} b \otimes_H 1 & \text{if } k = i + 1 \text{ and } b \text{ of type (II)} \\ r b \otimes_H 1 & \text{in all the other cases} \end{cases}$$

We note that the top equality is (CII') . Let's name the bottom equality (E) .

With Cases A, B, C, D, E , the action of the g_i 's on the vector space $B_1 \otimes_H F$ is entirely described. The object of the next part is to give an expression of the representation in terms of roots.

2.3 Expression of the representation in the Lawrence-Krammer space

2.3.1 The Lawrence-Krammer representation

Following our discussion at the end of § 1.3.2 and in § 2.1, there is a bijection:

$$u : \begin{array}{ccc} \phi^+ & \longrightarrow & B \\ \beta & \longmapsto & b \end{array},$$

where b is the algebra element associated with the positive root β , as in § 1.3.2. It follows that there is a natural isomorphism φ of vector spaces over F , defined on the basis vectors by:

$$\varphi : \begin{array}{ccc} \mathcal{V}^{(n)} & \xrightarrow{\sim} & B_1 \otimes_H F \\ x_\beta & \longmapsto & u(\beta) \otimes_H 1 \end{array}$$

We now get a representation of the BMW algebra inside the Lawrence-Krammer space as follows.

Theorem 3. *The map on the generators*

$$\nu^{(n)} : \begin{array}{ccc} B(A_{n-1}) & \longrightarrow & \text{End}_F(\mathcal{V}^{(n)}) \\ g_i & \longmapsto & \nu_i \end{array},$$

where each ν_i is defined on the basis vectors of $\mathcal{V}^{(n)}$ by

$$\nu_i(x_\beta) = \varphi^{-1}(g_i \cdot (u(\beta) \otimes_H 1))$$

defines a representation of degree $\frac{n(n-1)}{2}$ of the BMW algebra $B(A_{n-1})$ in the L-K space $\mathcal{V}^{(n)}$. Once irreducibility over $\mathbb{Q}(l, r)$ has been established, as a representation of the Braid group on n strands, it is equivalent to the Lawrence-Krammer representation.

Proof. By definition, $\nu^{(n)}$ is a representation of B in $\mathcal{V}^{(n)}$. We notice that $\nu^{(n)}$ factors through the quotient B/I_2 , where I_2 is the two sided ideal of B generated by all the products $e_i e_j$ with $|i - j| > 2$. Indeed, in B_1 , the algebra element $e_i e_j$ is zero and so in $B_1 \otimes_H F$, the vector $e_i e_j b \otimes_H 1$ is zero. Thus, we have:

$$\nu^{(n)}(e_i e_j) = 0 \text{ when } |i - j| > 2$$

Then by [2], as a representation of the Braid group on n strands, $\nu^{(n)}$ must be equivalent to the Lawrence-Krammer representation of the Artin group of type A_{n-1} based on the two parameters t and r , as defined in [3]. The r of this paper is the $\frac{1}{r}$ of [3]; the parameter t of [3] is related to the parameters l and r of this paper by $lt = r^3$. $\square$

2.3.2 An explicit form of the representation in terms of roots

Given a positive root

$$\beta = \alpha_i + \dots \alpha_j \text{ with } i < j,$$

we read on the expressions (AI) , (BI) , (CI) , (DI) and (E) of §2.2 an expression of $\nu_k(x_\beta)$ for $k \in \{i-1, i, j, j+1\}$ and for $k \notin \{i-1, i, j, j+1\}$. We define the height $ht(\beta)$ of a positive root β as the sum of its coefficients with respect to the simple roots $\alpha_1, \dots, \alpha_{n-1}$. We have:

$$\begin{aligned} \nu_{i-1}(x_\beta) &= x_{\beta+\alpha_{i-1}} + m r^{ht(\beta)-1} x_{\alpha_{i-1}} - m x_\beta && \text{by (AI)} \\ \nu_i(x_\beta) &= x_{\beta-\alpha_i} && \text{by (BI)} \\ \nu_j(x_\beta) &= x_{\beta-\alpha_j} + \frac{m}{l r^{ht(\beta)-2}} x_{\alpha_j} - m x_\beta && \text{by (CI)} \\ \nu_{j+1}(x_\beta) &= x_{\beta+\alpha_{j+1}} && \text{by (DI)} \\ \nu_k(x_\beta) &= r x_\beta \quad \forall k \notin \{i-1, i, j, j+1\} && \text{by (E)} \end{aligned}$$

Similarly for type (II) , if $\beta = \alpha_i$ is a simple root, we have:

$$\begin{aligned} \nu_{i-1}(x_\beta) &= x_{\beta+\alpha_{i-1}} + m x_{\alpha_{i-1}} - m x_{\alpha_i} && \text{by (AII)} \\ \nu_i(x_\beta) &= l^{-1} x_{\alpha_i} && \text{by (BII)} \\ \nu_{i+1}(x_\beta) &= x_{\beta+\alpha_{i+1}} && \text{by (CII')} \\ \nu_k(x_\beta) &= r x_\beta \quad \forall k \notin \{i-1, i, i+1\} \end{aligned}$$

The last equation is obtained with (CII) when $k > i + 1$ and with (E) when $k < i - 1$.

For each node i , we summarize the action of ν_i on x_β as follows.

$$\nu_i(x_\beta) = \begin{cases} r x_\beta & \text{if } (\beta|\alpha_i) = 0 \quad (a) \\ \frac{1}{l} x_\beta & \text{if } (\beta|\alpha_i) = 1 \quad (b) \\ x_{\beta+\alpha_i} & \text{if } (\beta|\alpha_i) = -\frac{1}{2} \quad \text{and (c)} \\ x_{\beta+\alpha_i} + m r^{ht(\beta)-1} x_{\alpha_i} - m x_\beta & \text{if } (\beta|\alpha_i) = -\frac{1}{2} \quad \text{and (c')} \\ x_{\beta-\alpha_i} + \frac{m}{l r^{ht(\beta)-2}} x_{\alpha_i} - m x_\beta & \text{if } (\beta|\alpha_i) = \frac{1}{2} \quad \text{and (d)} \\ x_{\beta-\alpha_i} & \text{if } (\beta|\alpha_i) = \frac{1}{2} \quad \text{and (d')} \end{cases}$$

where (c), (c'), (d), (d') are the following conditions:

$$\begin{aligned} (c) \quad \beta &= \alpha_t + \dots + \alpha_{i-1} & \text{with } t &\leq i-1 \\ (c') \quad \beta &= \alpha_{i+1} + \dots + \alpha_s & \text{with } s &\geq i+1 \\ (d) \quad \beta &= \alpha_t + \dots + \alpha_i & \text{with } t &\leq i-1 \\ (d') \quad \beta &= \alpha_i + \dots + \alpha_s & \text{with } s &\geq i+1 \end{aligned}$$

We note that:

(a) is equivalent to $Supp(\beta) \cap \{i-1, i, i+1\} = \emptyset$ or $\{i-1, i, i+1\} \subseteq Supp(\beta)$.

(b) is equivalent to $\beta = \alpha_i$.

(c) and (c') are the two ways the inner product $(\beta|\alpha_i)$ can be $-\frac{1}{2}$.

(d) and (d') are the two ways the inner product $(\beta|\alpha_i)$ can be $-\frac{1}{2}$.

We deduce from these equalities an expression for $\nu^{(n)}(e_i)$:

$$\nu^{(n)}(e_i)(x_\beta) = \begin{cases} 0 & \text{if } (\beta|\alpha_i) = 0 \\ \left(1 - \frac{l-\frac{1}{r}}{\frac{1}{r}-r}\right) x_{\alpha_i} & \text{if } (\beta|\alpha_i) = 1 \\ \frac{1}{r^{ht(\beta)-1}} x_{\alpha_i} & \text{if } (\beta|\alpha_i) = -\frac{1}{2} \quad \text{and (c)} \\ r^{ht(\beta)-1} x_{\alpha_i} & \text{if } (\beta|\alpha_i) = -\frac{1}{2} \quad \text{and (c')} \\ \frac{1}{l r^{ht(\beta)-2}} x_{\alpha_i} & \text{if } (\beta|\alpha_i) = \frac{1}{2} \quad \text{and (d)} \\ l r^{ht(\beta)-2} x_{\alpha_i} & \text{if } (\beta|\alpha_i) = \frac{1}{2} \quad \text{and (d')} \end{cases}$$

Notice $\nu^{(n)}(e_i)(x_\beta)$ is always a multiple of x_{α_i} . This is easily pictured on the tangles.

The next part establishes Theorem 1, following the discussion of §1.2.

3 Reducibility of the representation

3.1 Action on a proper invariant subspace of the L-K space

We show the following result:

Proposition 1.

For any proper invariant subspace $\mathcal{U}$ of $\mathcal{V}^{(n)}$, we have $\nu^{(n)}(e_i)(\mathcal{U}) = 0$ for all i .

Proof. If $\mathcal{U}$ is trivial, there is nothing to prove. Otherwise, let u be a nonzero vector of $\mathcal{U}$ such that $\nu^{(n)}(e_i)(u) \neq 0$. Since $\nu^{(n)}(e_i)(u)$ is a multiple of x_{α_i} , we see that x_{α_i} is in $\mathcal{U}$. From there, we have:

$$\nu_{i-1}(x_{\alpha_i}) = x_{\alpha_i + \alpha_{i-1}} + m x_{\alpha_{i-1}} \text{ modulo } F x_{\alpha_i}$$

Hence $x_{\alpha_i + \alpha_{i-1}} + m x_{\alpha_{i-1}}$ is in $\mathcal{U}$. Another application of ν_{i-1} now yields:

$$\nu_{i-1}(x_{\alpha_i + \alpha_{i-1}}) + m x_{\alpha_{i-1}} = x_{\alpha_i} + \frac{m}{l} x_{\alpha_{i-1}},$$

from which we derive that $x_{\alpha_{i-1}}$ is in $\mathcal{U}$. By induction, we see that all the x_{α_t} 's for $t \leq i$ are in $\mathcal{U}$. In particular, x_{α_1} is in $\mathcal{U}$. But since $e_1 \otimes_H 1$ spans $B_1 \otimes_H F$, x_{α_1} spans $\mathcal{V}^{(n)}$. Then $\mathcal{U}$ is the whole L-K space $\mathcal{V}^{(n)}$, in contradiction with our assumption that $\mathcal{U}$ is proper. $\square$

Corollary 1. *Let $\mathcal{W}$ be a proper irreducible invariant subspace of $\mathcal{V}^{(n)}$. Then, $\mathcal{W}$ is an irreducible $\mathcal{H}_{F,r^2}(n)$ -module.*

Proof. By Proposition 1 and (3), we have

$$[g_i^2 + m g_i - 1] \cdot \mathcal{W} = 0 \text{ for all } i.$$

Hence $\mathcal{W}$ is an $\mathcal{H}_{F,r^2}(n)$ -module. Since the e_i 's are polynomials in the g_i 's, $\mathcal{W}$ is an irreducible $\mathcal{H}_{F,r^2}(n)$ -module. $\square$

The next part investigates the existence of a one-dimensional invariant subspace of $\mathcal{V}^{(n)}$. We define for two nodes i and j with $i < j$

$$w_{ij} = x_{\alpha_i + \dots + \alpha_{j-1}}$$

We will sometimes write $w_{i,j}$ instead of w_{ij} . Below is how w_{ij} is represented in the tangle algebra:



It has two horizontal strands: one that joins nodes i and j at the top, and one that joins nodes 1 and 2 at the bottom and $(n - 2)$ vertical strands that don't cross within each other. The top horizontal strand over-crosses the vertical strands that it intersects.

3.2 The case $l = \frac{1}{r^{2n-3}}$

We will prove the theorem:

Theorem 4. *Let n be an integer with $n \geq 3$ and assume $(r^2)^2 \neq 1$.*

Suppose $n \geq 4$. There exists a one-dimensional invariant subspace of $\mathcal{V}^{(n)}$ if and only if $l = \frac{1}{r^{2n-3}}$. If so, it is spanned by $\sum_{1 \leq s < t \leq n} r^{s+t} w_{st}$

(Case $n = 3$) There exists a one-dimensional invariant subspace of $\mathcal{V}^{(3)}$ if and only

if $l = \frac{1}{r^3}$ or $l = -r^3$.

Moreover, if $r^6 \neq -1$, it is unique and

$$\begin{aligned} \text{when } l &= \frac{1}{r^3}, & \text{it is spanned by } & w_{12} + r w_{13} + r^2 w_{23} \\ \text{when } l &= -r^3, & \text{it is spanned by } & w_{12} - \frac{1}{r} w_{13} + \frac{1}{r^2} w_{23} \end{aligned}$$

If $r^6 = -1$, there are exactly two one-dimensional invariant subspaces of $\mathcal{V}^{(3)}$ and they are respectively spanned by the vectors above.

Proof. Let $\mathcal{U}$ be a one-dimensional invariant subspace of $\mathcal{V}^{(n)}$ and u a spanning vector of $\mathcal{U}$. For each i , let γ_i be the scalar such that $\nu_i(u) = \gamma_i u$. Since $(\nu_i^2 + m\nu_i - id_{\mathcal{V}^{(n)}})(u) = 0$ by Proposition 1, it follows that $\gamma_i^2 + m\gamma_i - 1 = 0$. Hence $\gamma_i \in \{r, -\frac{1}{r}\}$. Further, since $(r^2)^2 \neq 1$, the Braid relation $\nu_i\nu_j\nu_i = \nu_j\nu_i\nu_j$ when $i \sim j$ forces that γ_i takes the same value as γ_j . Let's denote by γ the common value of the γ_i 's. So, for each i , we have $\nu_i(u) = \gamma u$, where $\gamma \in \{r, -\frac{1}{r}\}$.

A general form for u is:

$$u = \sum_{1 \leq i < j \leq n} \mu_{ij} w_{ij}, \quad \text{where } \mu_{ij} \in F$$

We look for relations between these coefficients.

Lemma 3. *Let i be some node. Suppose $v = \sum_{1 \leq k < f \leq n} \mu_{kf} w_{kf}$ is a vector of $\mathcal{V}^{(n)}$ with $\nu_i(v) = \gamma v$ where $\gamma \in \{r, -\frac{1}{r}\}$. Then the following equalities hold for the coefficients of v :*

$$\forall s \geq i + 2, \mu_{i+1,s} = \gamma \mu_{i,s} \quad (14)$$

$$\forall t \leq i - 1, \mu_{t,i+1} = \gamma \mu_{t,i} \quad (15)$$

When $i = 1$, only (14) holds and when $i = n - 1$, only (15) holds.

Proof. To show (14), we look at the coefficient of $w_{i+1,s}$ in $\nu_i(v) = \gamma v$, where $s \geq i + 2$. We get: $\mu_{i,s} - m\mu_{i+1,s} = \gamma\mu_{i+1,s}$. Since $\gamma + m = \frac{1}{\gamma}$, this equality is equivalent to $\mu_{i+1,s} = \gamma\mu_{i,s}$. Similarly, by equating the coefficients of $w_{t,i+1}$ in $\nu_i(v) = \gamma v$, we obtain (15). $\square$

Applying these equalities to the coefficients of u , we see that all the coefficients of u must be nonzero. In particular, when $n \geq 4$, the coefficient μ_{34} of u is nonzero. Because an action of g_1 on w_{34} is a multiplication by r and an action of g_1 on the other terms w_{ij} does not create any term in w_{34} , this forces $\gamma = r$. Thus, without loss of generality, we have:

$$u = \sum_{1 \leq i < j \leq n} r^{i+j} w_{ij}$$

From there, let's look at the action of g_1 on u and the resulting coefficient in w_{12} . The action of g_1 on w_{12} is a multiplication by l^{-1} and an action of g_1 on the $w_{2,j}$'s for $3 \leq j \leq n$ creates new terms in w_{12} with respective coefficients $m r^{j-3}$. Thus, we get the equation:

$$\frac{r^3}{l} + \sum_{j=3}^n (r^2)^j = r^4,$$

from which we derive that $l = \frac{1}{r^{2n-3}}$.

Conversely, if $l = \frac{1}{r^{2n-3}}$, we define u as $\sum_{1 \leq i < j \leq n} r^{i+j} w_{ij}$ and check that $\nu_i(u) = r u$ for each i . First we show that the coefficient r^{2i+1} of $w_{i,i+1}$ is multiplied by r when acting by g_i . There are three contributions. One comes from the terms $w_{t,i+1}$, with coefficient $r^{t+i+1} \frac{m r^{2n-3}}{r^{i-t-1}}$. Another one from the terms $w_{i+1,s}$ with coefficient $r^{s+i+1} m r^{s-i-2}$, and a third one simply from the term $w_{i,i+1}$, with coefficient $r^{2i+1} r^{2n-3}$. Now, we have:

$$m r^{2n-1} \sum_{t=1}^{i-1} r^{2t} + \frac{m}{r} \sum_{s=i+2}^n r^{2s} + r^{2n+2i-2} = r r^{2i+1}$$

Next, given a positive root β , if none of the nodes $i-1$, i , $i+1$ is in the support of β or if all three nodes $i-1$, i , $i+1$ are in the support of β , then it comes:

$$\nu_i(x_\beta) = r x_\beta$$

Thus, we only need to study the action of ν_i on $w_{k,i}$, $w_{k,i+1}$, with $k \leq i-1$ on one hand and $w_{i,l}$, $w_{i+1,l}$, with $l \geq i+2$ on the other hand.

We have:

$$\begin{aligned} r^{k+i} \nu_i(w_{k,i}) &= r^{k+i} w_{k,i+1} \\ r^{k+i+1} \nu_i(w_{k,i+1}) &= r^{k+i+1} w_{k,i} - m r^{k+i+1} w_{k,i+1} \text{ modulo } F x_{\alpha_i} \end{aligned}$$

So we get:

$$\nu_i(r^{k+i} w_{k,i} + r^{k+i+1} w_{k,i+1}) = r^{k+i+1} w_{k,i} + r^{k+i+2} w_{k,i+1} \text{ modulo } F x_{\alpha_i}$$

Similarly, we have:

$$\begin{aligned} r^{l+i} \nu_i(w_{i,l}) &= r^{l+i} w_{i+1,l} \\ r^{l+i+1} \nu_i(w_{i+1,l}) &= r^{l+i+1} w_{i,l} - m r^{l+i+1} w_{i+1,l} \text{ modulo } F x_{\alpha_i}, \end{aligned}$$

so that:

$$\nu_i(r^{l+i} w_{i,l} + r^{l+i+1} w_{i+1,l}) = r^{l+i+1} w_{i,l} + r^{l+i+2} w_{i+1,l} \text{ modulo } F x_{\alpha_i}$$

This ends the proof of the Theorem when $n \geq 4$.

Suppose now $n = 3$. So,

$$u = \gamma^3 w_{12} + \gamma^4 w_{13} + \gamma^5 w_{23}$$

Let's compute $\nu_1(u)$ and $\nu_2(u)$:

$$\begin{aligned} \nu_1(u) &= \left(\frac{\gamma^3}{l} + m \gamma^5\right) w_{12} + \gamma^5 w_{13} + \gamma^6 w_{23} \\ \nu_2(u) &= \gamma^4 w_{12} + \gamma^5 w_{13} + \left(\frac{\gamma^5}{l} + \frac{m \gamma^4}{l}\right) w_{23} \end{aligned}$$

Since $\nu_1(u) = \gamma u$, we must have:

$$\frac{\gamma^3}{l} + m \gamma^5 = \gamma^4, \quad \text{i.e.} \quad \frac{1}{l} = \gamma(1 - m \gamma), \quad \text{i.e.} \quad l = \frac{1}{\gamma^3}, \quad \text{as} \quad 1 - m \gamma = \gamma^2$$

Thus, if there exists a one dimensional invariant subspace of $\mathcal{V}^{(3)}$ then l must take the values $\frac{1}{r^3}$ or $-r^3$.

Conversely, let's consider the two vectors:

$$\begin{aligned} u_r &= w_{12} + r w_{13} + r^2 w_{23} \\ u_{-\frac{1}{r}} &= w_{12} - \frac{1}{r} w_{13} + \frac{1}{r^2} w_{23} \end{aligned}$$

We read on the equations giving the expressions for $\nu_1(u)$ and $\nu_2(u)$ that

$$\begin{aligned} \text{If } l &= \frac{1}{r^3} \text{ then } \nu_1(u_r) = \nu_2(u_r) = r u_r \\ \text{If } l &= -r^3 \text{ then } \nu_1(u_{-\frac{1}{r}}) = \nu_2(u_{-\frac{1}{r}}) = -\frac{1}{r} u_{-\frac{1}{r}} \quad \square \end{aligned}$$

The next section investigates the existence of an irreducible $(n-1)$ -dimensional invariant subspace of $\mathcal{V}^{(n)}$.

3.3 The case $l \in \{\frac{1}{r^{n-3}}, -\frac{1}{r^{n-3}}\}$

In Theorem 4, the case $n = 3$ was special. Likewise, in the following Theorem 5, the case $n = 4$ needs to be formulated separately.

Theorem 5.

Let n be a positive integer with $n \geq 3$ and $n \neq 4$. Assume $\mathcal{H}_{F,r^2}(n)$ is semisimple. Then, there exists an irreducible $(n-1)$ -dimensional invariant subspace of $\mathcal{V}^{(n)}$ if and only if $l = \frac{1}{r^{n-3}}$ or $l = -\frac{1}{r^{n-3}}$.

If so, it is spanned by the $v_i^{(n)}$'s, $1 \leq i \leq n-1$, where $v_i^{(n)}$ is defined by the formula:

$$\begin{aligned} v_i^{(n)} &= \left(\frac{1}{r} - \frac{1}{l}\right) w_{i,i+1} + \sum_{s=i+2}^n r^{s-i-2} (w_{i,s} - \frac{1}{r} w_{i+1,s}) \\ &\quad + \epsilon_l \sum_{t=1}^{i-1} r^{n-i-2+t} (w_{t,i} - \frac{1}{r} w_{t,i+1}) \end{aligned}$$

$$\text{with } \begin{cases} \epsilon_{\frac{1}{r^{n-3}}} = 1 \\ \epsilon_{-\frac{1}{r^{n-3}}} = -1 \end{cases}$$

(Case $n = 4$) Assume $\mathcal{H}_{F,r^2}(4)$ is semisimple. Then, there exists an irreducible 3-dimensional invariant subspace of $\mathcal{V}^{(4)}$ if and only if $l \in \{\frac{1}{r}, -\frac{1}{r}, -r^3\}$.

If $l \in \{-\frac{1}{r}, \frac{1}{r}\}$, it is spanned by $v_1^{(4)}, v_2^{(4)}, v_3^{(4)}$

If $l = -r^3$, it is spanned by the vectors:

$$\begin{cases} u_1 &= r w_{23} + w_{13} + (\frac{1}{r} + \frac{1}{r^3}) w_{34} - w_{24} - \frac{1}{r} w_{14} \\ u_2 &= -r w_{12} - r^2 w_{13} - \frac{1}{r} w_{34} - \frac{1}{r^2} w_{24} + (r + \frac{1}{r}) w_{14} \\ u_3 &= (r + \frac{1}{r^3}) w_{12} + \frac{1}{r} w_{23} - w_{13} + w_{24} - r w_{14} \end{cases}$$

Proof. Suppose that there exists an irreducible $(n-1)$ -dimensional invariant subspace $\mathcal{U}$ of $\mathcal{V}^{(n)}$.

Claim 1. *Except in the case when $n = 6$, there exists a basis $(v_1, \dots, v_{n-1})$ of $\mathcal{U}$ such that one of the following two sets of relations holds:*

$$\begin{aligned}
 (\triangle) \quad & \begin{cases} \nu_t(v_i) = r v_i & \forall t \notin \{i-1, i, i+1\} \\ \nu_i(v_i) = -\frac{1}{r} v_i & \forall 1 \leq i \leq n-1 \\ \nu_{i+1}(v_i) = r(v_i + v_{i+1}) & \forall 1 \leq i \leq n-2 \\ \nu_{i-1}(v_i) = r v_i + \frac{1}{r} v_{i-1} & \forall 2 \leq i \leq n-1 \end{cases} \\
 (\nabla) \quad & \begin{cases} \nu_t(v_i) = -1/r v_i & \forall t \notin \{i-1, i, i+1\} \\ \nu_i(v_i) = r v_i & \forall 1 \leq i \leq n-1 \\ \nu_{i+1}(v_i) = -1/r(v_i + v_{i+1}) & \forall 1 \leq i \leq n-2 \\ \nu_{i-1}(v_i) = -1/r v_i - r v_{i-1} & \forall 2 \leq i \leq n-1 \end{cases}
 \end{aligned}$$

Proof. We first recall some general fact about the irreducible representations of the Iwahori-Hecke algebra of the symmetric group. The following result was established by James for the irreducible representations of the symmetric group, but applies here to the Iwahori-Hecke algebra $\mathcal{H}_{F,r^2}(n)$ since we work in characteristic zero and assumed $\mathcal{H}_{F,r^2}(n)$ semisimple. By Theorem 6, point (i) in [4], when the characteristic of the field F is zero and for $n \geq 7$, an irreducible $F \text{Sym}(n)$ -module is either one of the Specht modules $S^{(n)}$, $S^{(1^n)}$, $S^{(n-1,1)}$, $S^{(2,1^{n-2})}$ or has dimension greater than $(n-1)$. The statement is also true when $n = 3$ and $n = 5$. When $n = 4$, the statement does not hold as $S^{(2,2)}$ has dimension 2 and when $n = 6$, the statement also fails since $S^{(3,3)}$ and $S^{(2,2,2)}$ both have dimension 5. In any case, there are exactly two inequivalent irreducible representations of $F \text{Sym}(n)$ of degree $(n-1)$, except in the case $n = 6$, when there are exactly four inequivalent irreducible representations of $F \text{Sym}(6)$ of degree 5. The same statement holds for $\mathcal{H}_{F,r^2}(n)$ when the algebra is semisimple.

Consider now the set of relations $(\triangle)$ (resp (∇)). For each i , let M_i (resp N_i) be the matrix of the endomorphism ν_i in the basis $(v_1, \dots, v_{n-1})$. It is a direct verification that the M_i 's (resp N_i 's) satisfy the Braid relations and the relation $M_i^2 + m M_i = I_{n-1}$ (resp $N_i^2 + m N_i = I_{n-1}$) for each i , where I_{n-1} is the identity matrix of size $(n-1)$. Hence the M_i 's (resp the N_i 's) yield a matrix representation of $\mathcal{H}_{F,r^2}(n)$ of degree $(n-1)$.

To show that these two matrix representations are irreducible, relying on James's statement above, it suffices to check that there is no one-dimensional invariant subspace of F^{n-1} when $n \neq 4$ and that there is no one-dimensional or irreducible two-dimensional invariant subspace of F^3 when $n = 4$. This is the case if $r^{2n} \neq 1$ when $n \neq 4$ and if $(r^2)^2 \neq 1$ and $(r^2)^4 \neq 1$ when $n = 4$.

When $n = 3$, the two matrix representations are equivalent. When $n \geq 4$, they are not: visibly, the matrices of one representation all have the same trace $-\frac{(n-2)}{r} + r$ and the matrices of the other one all have the same trace $(n-2)r - \frac{1}{r}$. These two values are distinct when $(r^2)^2 \neq 1$ and $n \geq 4$. We conclude that these are the two inequivalent irreducible representations of $\mathcal{H}_{F,r^2}(n)$ when $n \geq 4$ and $n \neq 6$. $\square$

In what follows, we assume $n \geq 4$. We will show that it is impossible to have the second set of relations, except in the case $n = 4$ when it forces $l = -r^3$. Suppose the v_i 's satisfy (∇) . The relation $\nu_{n-1}(v_1) = -\frac{1}{r} v_1$ implies that in v_1 there are no terms in $w_{s,t}$ for integers $s, t \in \{1, \dots, n-2\}$ such that $s < t$.

Hence we may write:

$$v_1 = \sum_{j=1}^{n-2} \mu_{j,n-1} w_{j,n-1} + \sum_{j=1}^{n-1} \mu_{j,n} w_{j,n}$$

Moreover, the relation $\nu_3(v_1) = -\frac{1}{r} v_1$ implies that there are no terms in $w_{j,k}$ in v_1 for $j \geq 5$. Further, the relations

$$\begin{cases} \nu_1(w_{2,n-1}) &= w_{1,n-1} + mr^{n-4} w_{12} - m w_{2,n-1} \\ \nu_1(w_{2,n}) &= w_{1,n} + mr^{n-3} w_{12} - m w_{2,n} \end{cases}$$

imply that: $mr^{n-4} \mu_{2,n-1} + mr^{n-3} \mu_{2,n} = 0$, i.e. $\mu_{2,n} = -\frac{1}{r} \mu_{2,n-1}$, as there is no term in w_{12} in v_1 . Furthermore, an application of (14) with $\gamma = r$ and $i = 1$ yields for $s = n-1$ and $s = n$ respectively: $\mu_{2,n-1} = r \mu_{1,n-1}$ and $\mu_{2,n} = r \mu_{1,n}$. So, up to a multiplication by a scalar,

$$\begin{aligned} v_1 &= w_{1,n-1} + r w_{2,n-1} - \frac{1}{r} w_{1,n} - w_{2,n} \\ &\quad + \mu_{3,n-1} w_{3,n-1} + \mu_{4,n-1} w_{4,n-1} + \mu_{3,n} w_{3,n} + \mu_{4,n} w_{4,n} \end{aligned} \quad (16)$$

$$\text{or} \quad v_1 = \mu_{3,n-1} w_{3,n-1} + \mu_{4,n-1} w_{4,n-1} + \mu_{3,n} w_{3,n} + \mu_{4,n} w_{4,n} \quad (17)$$

If $n \geq 5$, the relation $\nu_3(v_1) = -\frac{1}{r} v_1$ implies that $\mu_{2,n} = 0$. Then an expression for v_1 is given by (17) and not (16). Assume $n > 5$. Then there is no term in w_{34} in v_1 . Then it comes $\mu_{4,n} = -\frac{1}{r} \mu_{4,n-1}$. Moreover, by (14) applied with $\gamma = -\frac{1}{r}$ and $i = 3$, we have $\mu_{4,n} = -\frac{1}{r} \mu_{3,n}$ and $\mu_{4,n-1} = -\frac{1}{r} \mu_{3,n-1}$. Further, when $n > 5$, $\nu_4(w_{3,n}) = r w_{3,n}$ and an action of g_4 on the other terms of v_1 in (17) won't create any term in $w_{3,n}$. Thus, the relation $\nu_4(v_1) = -\frac{1}{r} v_1$ forces $\mu_{3,n} = 0$. Then, by the relations previously established, all the coefficients of v_1 are actually zero, which is impossible. The case $n = 5$ also leads to a contradiction and details appear in [7], §8.3.

Suppose now $n = 4$. By (16) and (17), $v_1 = w_{13} + r w_{23} - \frac{1}{r} w_{14} - w_{24} + \mu_{34} w_{34}$ or $v_1 = w_{34}$. Suppose v_1 is of the second type. Then, by $\nu_3(v_1) = -\frac{1}{r} v_1$, we must have $l = -r$. Since $\nu_2(v_1) = -\frac{1}{r}(v_1 + v_2)$, we must have:

$$v_2 = -r w_{24} + (r^2 - 1) w_{23} - r^2 w_{34}$$

Since $\begin{cases} \nu_2(w_{34}) = w_{24} + m w_{23} - m w_{34} \\ \nu_2(w_{23}) = -\frac{1}{r} w_{23} \end{cases}$ and since $\nu_2(v_2) = r v_2$, we get:

$$-m r^2 - \frac{1}{r}(r^2 - 1) = r(r^2 - 1), \text{ which reads } m = 0 \text{ after simplification.}$$

As m is nonzero, this is a contradiction. Thus, v_1 is of the first type. Then, denoting by λ_{ij} the coefficient of w_{ij} in v_2 , we get by looking at the coefficient of w_{12} in $\nu_2(v_1) = -\frac{1}{r} v_1 - \frac{1}{r} v_2$ that $\lambda_{12} = -r$. Since by (15) with $\gamma = r$ and $i = 2$, we have $\lambda_{13} = r w_{12}$, it follows that $\lambda_{13} = -r^2$. Next, by looking at the coefficient of w_{14} in the relation $\nu_2(v_1) = -\frac{1}{r} v_1 - \frac{1}{r} v_2$, we get:

$$-1 = -\frac{\lambda_{14}}{r} + \frac{1}{r^2} \quad \text{i.e.} \quad \lambda_{14} = r + \frac{1}{r}$$

Also, by looking at the coefficient of w_{24} in the same relation, we obtain:

$$\mu_{34} = \frac{1}{r} - \frac{\lambda_{24}}{r} \quad (18)$$

Next, we use the relation $\nu_1(v_2) = -\frac{1}{r}v_2 - r v_1$. First we look at the coefficient of w_{13} to get $\lambda_{23} = 0$ and by looking at the coefficient of w_{14} , we get $\lambda_{24} = -\frac{1}{r^2}$. By $\nu_2(v_2) = r v_2$ and (14), it comes $\lambda_{34} = r \lambda_{24} = -\frac{1}{r}$. Plugging the value of λ_{24} in (18) now yields $\mu_{34} = \frac{1}{r} + \frac{1}{r^3}$. Finally, by looking at the coefficient of w_{12} in $\nu_1(v_2) = -\frac{1}{r}v_2 - r v_1$, we get $-\frac{r}{l} - \frac{m}{r} = 1$, from which we derive $l = -r^3$. Also, gathering all the results above, we see that the vectors v_1 and v_2 are exactly the vectors u_1 and u_2 of the Theorem. Similar computations would also lead to $v_3 = u_3$ (see [7]). Conversely, we can show that if $l = -r^3$, then the vectors u_1, u_2 and u_3 form a free family of vectors that satisfy the relations (∇) . This shows that their linear span over F is an irreducible 3-dimensional invariant subspace of $\mathcal{V}^{(4)}$. For details, see [7], §8.3.

Suppose now that the v_i 's satisfy (Δ) . The relation $\nu_i(v_i) = -\frac{1}{r}v_i$ implies that in v_i there are no terms in w_{ts} for $s \leq i-1$ or $t \geq i+2$ or $t \leq i-1$ and $s \geq i+2$. Thus, a general form for v_i must be:

$$\begin{aligned} v_i = & \mu_{i,i+1} w_{i,i+1} + \sum_{s=i+2}^n \mu_{i,s} w_{i,s} + \sum_{s=i+2}^n \mu_{i+1,s} w_{i+1,s} \\ & + \sum_{t=1}^{i-1} \mu_{t,i} w_{t,i} + \sum_{t=1}^{i-1} \mu_{t,i+1} w_{t,i+1} \end{aligned} \quad (19)$$

Since $\nu_i(v_i) = -\frac{1}{r}v_i$, both equalities (14) and (15) hold with $\gamma = -\frac{1}{r}$. Further, since $\nu_q(v_i) = r v_i$ for $q \notin \{i-1, i, i+1\}$, applying (14) and (15) with $i = q$ and $\gamma = r$ yields:

$$\forall j \geq q+2, \mu_{q+1,j} = r, \mu_{q,j} \quad (20)$$

$$\forall k \leq q-1, \mu_{k,q+1} = r \mu_{k,q} \quad (21)$$

Apply (20) with $q \leq i-2$ and $j \in \{i, i+1\}$ to get:

$$\forall q \leq i-2, \mu_{q+1,i} = r \mu_{q,i} \quad \& \quad \mu_{q+1,i+1} = r \mu_{q,i+1}$$

Apply (21) with $q \geq i+2$ and $k \in \{i, i+1\}$ to get:

$$\forall q \geq i+2, \mu_{i,q+1} = r \mu_{i,q} \quad \& \quad \mu_{i+1,q+1} = r \mu_{i+1,q}$$

Expression (19) may now be rewritten:

$$v_i = \zeta^{(i)} w_{i,i+1} + \delta^{(i)} \sum_{s=i+2}^n r^{s-i-2} (w_{i,s} - \frac{1}{r} w_{i+1,s}) + \lambda^{(i)} \sum_{t=1}^{i-1} r^{t-1} (w_{t,i} - \frac{1}{r} w_{t,i+1}),$$

where $\zeta^{(i)}$, $\delta^{(i)}$ and $\lambda^{(i)}$ are three coefficients to determine. First, we show that all the $\delta^{(i)}$ with $i \in \{1, \dots, n-2\}$ may be set to the value one. Notice that if $v_1, \dots, v_{n-1}$ satisfy (Δ) , then $\delta v_1, \dots, \delta v_{n-1}$ also satisfy (Δ) , where δ is any nonzero scalar. Then, without loss of generality, we set $\delta^{(1)} = 1$. Suppose $\delta^{(i)} = 1$ for some node i with $1 \leq i \leq n-2$. We will show that $\delta^{(i+1)} = 1$.

Notice that $\delta^{(i+1)}$ is the coefficient of $w_{i+1,i+3}$ in v_{i+1} . Since an action of g_{i+1} on v_i never creates a term in $w_{i+1,i+3}$, by looking at the coefficient of $w_{i+1,i+3}$ in $\nu_{i+1}(v_i) = r v_i + r v_{i+1}$, we get $0 = -r \delta^{(i)} + r \delta^{(i+1)}$. After replacing $\delta^{(i)}$ by 1, this yields $\delta^{(i+1)} = 1$. Thus, all the $\delta^{(i)}$ may be set to the value 1. It remains to find the coefficients $\zeta^{(i)}$ and $\lambda^{(i)}$. By looking at the coefficient of $w_{i,i+1}$ in $\nu_{i+1}(v_i) = r(v_i + v_{i+1})$, we get:

$$r \zeta^{(i)} + r^i \lambda^{(i+1)} = 1, \text{ for each } i \text{ with } 1 \leq i \leq n-2 \quad (22)$$

Also, by looking at the coefficient of the same term $w_{i,i+1}$ in the relation $\nu_{i-1}(v_i) = r v_i + \frac{1}{r} v_{i-1}$, we get:

$$-m \zeta^{(i)} - r^{i-3} \lambda^{(i)} = r \zeta^{(i)} - \frac{1}{r^2}, \text{ for each } i \text{ with } 2 \leq i \leq n-1$$

After multiplication by a factor r^2 , we obtain:

$$r \zeta^{(i)} + r^{i-1} \lambda^{(i)} = 1, \text{ for each } i \text{ with } 2 \leq i \leq n-1 \quad (23)$$

By (22) and (23), we get $\lambda^{(i)} = \frac{1}{r^{i-2}} \lambda^{(2)}$, for all $i \geq 2$. Let's do a change of indices in (22) to get:

$$r \zeta^{(i-1)} + r^{i-1} \lambda^{(i)} = 1 \text{ for each } i \text{ with } 2 \leq i \leq n-1 \quad (24)$$

(23) and (24) show that $\zeta^{(i)} = \zeta^{(i-1)}$ for each i with $2 \leq i \leq n-1$. In other words, all the $\zeta^{(i)}$ are equal to a certain scalar ζ . The relation between ζ and $\lambda^{(2)}$ is given by equation (24) with $i = 2$:

$$\lambda^{(2)} = \frac{1}{r} - \zeta \quad (25)$$

Thus, by determining ζ , we will get a complete expression for all the vectors v_i 's. Since we have

$$v_1 = \zeta w_{12} + \sum_{s=3}^n r^{s-3} (w_{1,s} - \frac{1}{r} w_{2,s}),$$

by looking at the coefficient of w_{12} in the relation $\nu_1(v_1) = -\frac{1}{r} v_1$, we get the equation:

$$\zeta \left(\frac{1}{l} + \frac{1}{r} \right) = \frac{1}{r^2} - (r^2)^{n-3} \quad (26)$$

Further, by looking at the coefficient of $w_{i,i+1}$ in $\nu_i(v_i) = -\frac{1}{r} v_i$, we have:

$$\zeta \left(\frac{1}{l} + \frac{1}{r} \right) = \sum_{s=i+2}^n r^{s-i-3} m r^{s-i-2} + \lambda^{(i)} \sum_{t=1}^{i-1} r^{t-2} \frac{m}{l r^{i-t-1}}$$

i.e

$$\zeta \left(\frac{1}{l} + \frac{1}{r} \right) = \frac{1}{r^2} - (r^2)^{n-i-2} + \frac{\lambda^{(i)}}{l} \left(\frac{1}{r^i} - r^{i-2} \right) \quad (\star)_i$$

Let's write down $(\star)_2$ and $(\star)_3$:

$$\zeta \left(\frac{1}{l} + \frac{1}{r} \right) = \frac{1}{r^2} - (r^2)^{n-4} + \frac{\lambda^{(2)}}{l} \left(\frac{1}{r^2} - 1 \right) \quad (\star)_2$$

$$\zeta \left(\frac{1}{l} + \frac{1}{r} \right) = \frac{1}{r^2} - (r^2)^{n-5} + \frac{\lambda^{(2)}}{l r} \left(\frac{1}{r^3} - r \right) \quad (\star)_3$$

where $\lambda^{(3)}$ has been replaced by $\frac{\lambda^{(2)}}{r}$. Let's subtract these two equalities:

$$\frac{\lambda^{(2)}}{l} \left(\frac{1}{r^2} - \frac{1}{r^4} \right) = (r^2)^{n-4} \left(1 - \frac{1}{r^2} \right) \quad (\star)_2 - (\star)_3$$

After multiplying this equality by $\frac{1}{r^2}$ and dividing it by $\frac{1}{r^2} - \frac{1}{r^4}$ (licit as $m \neq 0$), we obtain:

$$\lambda^{(2)} = l (r^2)^{n-3} \quad (27)$$

Hence, by (25), $\zeta = \frac{1}{r} - l (r^2)^{n-3}$. Plugging this value for ζ into (26) now yields:

$$l^2 = \frac{1}{(r^2)^{n-3}}, \text{ hence } l \in \left\{ \frac{1}{r^{n-3}}, -\frac{1}{r^{n-3}} \right\}$$

If $l = \frac{1}{r^{n-3}}$, we get successively $\lambda^2 = r^{n-3} = \frac{1}{l}$, $\zeta = \frac{1}{r} - \frac{1}{l}$ and $\lambda^i = r^{n-i-1}$.

If $l = -\frac{1}{r^{n-3}}$, λ^2 and ζ are still respectively $\frac{1}{l}$ and $\frac{1}{r} - \frac{1}{l}$ and $\lambda^i = -r^{n-i-1}$. We obtain the formula announced in Theorem 5.

Conversely, if $l \in \left\{ \frac{1}{r^{n-3}}, -\frac{1}{r^{n-3}} \right\}$, we can show that the $v_i^{(n)}$'s defined in Theorem 5 satisfy the relations $(\triangle)$ (see [7], §8.3). In particular, their linear span over F is a proper invariant subspace of $\mathcal{V}^{(n)}$, hence is an $\mathcal{H}_{F,r^2}(n)$ -module by Corollary 1. When $n \neq 4$, if the vectors $v_i^{(n)}$'s were linearly dependent, then their span would either be one-dimensional or would contain a one-dimensional $\mathcal{H}_{F,r^2}(n)$ -submodule, as there is no irreducible $\mathcal{H}_{F,r^2}(n)$ -module of dimension between 1 and $(n-1)$. In any case, by Theorem 4, that would force $l = \frac{1}{r^{2n-3}}$ when $n \neq 3$ and $l \in \{-r^3, \frac{1}{r^3}\}$ when $n = 3$. This is impossible with our assumption that $l \in \left\{ \frac{1}{r^{n-3}}, -\frac{1}{r^{n-3}} \right\}$ and the fact that $r^{2n} \neq 1$. As for $n = 4$, the freedom over F of the family of vectors $(v_1^{(4)}, v_2^{(4)}, v_3^{(4)})$ is a direct verification or is a consequence of Theorem 4 and forthcoming Proposition 3 (See §3.4). We are now able to conclude: the vector space $\text{Span}_F(v_1^{(n)}, \dots, v_{n-1}^{(n)})$ is $(n-1)$ -dimensional, is invariant under the action of the g_i 's and is an $\mathcal{H}_{F,r^2}(n)$ -module since it is a proper invariant subspace of $\mathcal{V}^{(n)}$. Then, by the relations satisfied by the $v_i^{(n)}$'s, it must be irreducible.

To complete the proof of Theorem 5, we show that there does not exist any irreducible 5-dimensional invariant subspace of $\mathcal{V}^{(6)}$ that is isomorphic to one of the Specht modules $S^{(3,3)}$ or $S^{(2,2,2)}$. Indeed, suppose such a subspace exists and name it $\mathcal{W}$. Since we have assumed that $\mathcal{H}_{F,r^2}(6)$ is semisimple, it is licit to use the branching rule as it is described in Corollary 6.2 of [8]. We have:

$$S^{(3,3)} \downarrow_{\mathcal{H}_{F,r^2}(5)} \simeq S^{(3,2)} \quad S^{(2,2,2)} \downarrow_{\mathcal{H}_{F,r^2}(5)} \simeq S^{(2,2,1)}$$

We will show that the restriction of $\mathcal{W}$ to $\mathcal{H}_{F,r^2}(5)$ cannot be isomorphic to $S^{(3,2)}$ or $S^{(2,2,1)}$, hence a contradiction. A proof of the following fact is in [7], §8.3

Fact 1. Suppose $\mathcal{H}_{F,r^2}(5)$ is semisimple. Then, up to equivalence, the two irreducible matrix representations of degree 5 of $\mathcal{H}_{F,r^2}(5)$ are respectively defined by the matrices P_1, P_2, P_3, P_4 and Q_1, Q_2, Q_3, Q_4 given by:

$$P_1 := \begin{bmatrix} r & & & & \\ & r & & & \\ & & r & & \\ 1 & & -r^2 & -\frac{1}{r} & \\ & 1 & & -\frac{1}{r} & \end{bmatrix}, \quad P_2 := \begin{bmatrix} -\frac{1}{r} & & & 1 & \\ & -\frac{1}{r} & 1 & & \\ & & r & & \\ & & & r & \\ & & & & r \end{bmatrix}$$

$$P_3 := \begin{bmatrix} r & & & & \\ & r & & & \\ & 1 & -\frac{1}{r} & & \\ 1 & & & -\frac{1}{r} & -r^2 \\ & & & r & \end{bmatrix}, P_4 := \begin{bmatrix} & 1 & -r & & \\ 1 & r - \frac{1}{r} & 1 & & \\ & r & -r^2 & 1 & \\ & & r & 1 & r - \frac{1}{r} \end{bmatrix}$$

and for the conjugate representation:

$$Q_1 := \begin{bmatrix} -\frac{1}{r} & & & & \\ & -\frac{1}{r} & & & \\ & & -\frac{1}{r} & & \\ 1 & & -\frac{1}{r^2} & r & \\ & 1 & & & r \end{bmatrix}, Q_2 := \begin{bmatrix} r & & 1 & & \\ & r & -\frac{1}{r} & & \\ & & -\frac{1}{r} & & \\ & & & -\frac{1}{r} & \\ & & & & -\frac{1}{r} \end{bmatrix}$$

$$Q_3 := \begin{bmatrix} -\frac{1}{r} & & & & \\ & -\frac{1}{r} & & & \\ & 1 & r & & \\ 1 & & & r & -\frac{1}{r^2} \\ & & & -\frac{1}{r} & \end{bmatrix}, Q_4 := \begin{bmatrix} & 1 & \frac{1}{r} & & \\ 1 & r - \frac{1}{r} & -\frac{1}{r^2} & & \\ & & -\frac{1}{r} & & \\ & & -\frac{1}{r} & 1 & r - \frac{1}{r} \end{bmatrix}$$

where the blanks must be filled with zeros.

First we show that it is impossible to have a basis $(w_1, w_2, w_3, w_4, w_5)$ of $\mathcal{W}$ in which the matrices of the left action by the g_i 's, $i = 1, \dots, 4$ are the Q_i 's. Indeed, suppose that such a basis of vectors exists. Let's denote by $\lambda_{ij}^{(k)}$ the coefficient of w_{ij} in w_k . Since $g_4.w_4 = w_5$ and $g_3.w_5 = -\frac{1}{r^2}w_4 - \frac{1}{r}w_5$, we get

$$g_3g_4w_4 = -\frac{1}{r}g_4.w_4 - \frac{1}{r^2}w_4$$

We look at the coefficient of w_{12} in this equation to get

$$r^2 \lambda_{12}^{(4)} = -\lambda_{12}^{(4)} - \frac{1}{r^2} \lambda_{12}^{(4)}$$

Since $(r^2)^3 \neq 0$, we have $r^2 + 1 + \frac{1}{r^2} \neq 0$ and so $\lambda_{12}^{(4)} = 0$. Now since $g_3.w_1 = -\frac{1}{r}w_1 + w_4$, we get $r \lambda_{12}^{(1)} = -\frac{1}{r} \lambda_{12}^{(1)}$ and so $\lambda_{12}^{(1)} = 0$. This implies that $\lambda_{13}^{(1)}$ is also zero by $g_2.w_1 = r w_1$ and Lemma 3. Then, by looking at the coefficient of w_{13} in $g_2.w_4 = w_1 - \frac{1}{r}w_4$, we get $-m \lambda_{13}^{(4)} = -\frac{1}{r} \lambda_{13}^{(4)}$, where we used that $\lambda_{12}^{(4)} = 0$. Thus, $\lambda_{13}^{(4)} = 0$. Since $g_3.w_4 = r w_4$ and $g_1.w_4 = r w_4$, by Lemma 3, we also get:

$$\lambda_{13}^{(4)} = \lambda_{23}^{(4)} = \lambda_{14}^{(4)} = \lambda_{24}^{(4)} = 0$$

Let's now look at the term in w_{25} in $g_3g_4w_4 = -\frac{1}{r}g_4.w_4 - \frac{1}{r^2}w_4$. We have:

$$-mr^2 \lambda_{15}^{(4)} = -\frac{1}{r} \left(-mr \lambda_{15}^{(4)} \right) - \frac{1}{r^2} r \lambda_{15}^{(4)},$$

where we used that $\lambda_{25}^{(4)} = r \lambda_{15}^{(4)}$. Then, $\lambda_{15}^{(4)} = 0$ and also $\lambda_{25}^{(4)} = 0$. Further, since $\lambda_{12}^{(4)} = 0$, in $g_1.w_4$, a term in w_{12} is created only when g_1 acts on w_{26} , with coefficient $m r^3$. Thus the relation $g_1.w_4 = r w_4$ yields $\lambda_{26}^{(4)} = 0$. Then by $g_1.w_4 = r w_4$, we also have $\lambda_{16}^{(4)} = 0$.

Furthermore, on one hand, by looking at the coefficient of w_{34} in $g_1.w_1 = -\frac{1}{r}w_1 + w_4$, we get $r\lambda_{34}^{(1)} = -\frac{1}{r}\lambda_{34}^{(1)} + \lambda_{34}^{(4)}$, i.e

$$\lambda_{34}^{(4)} = \left(r + \frac{1}{r}\right)\lambda_{34}^{(1)}$$

On the other hand, by looking at the coefficient of w_{34} in $g_2.w_4 = w_1 - \frac{1}{r}w_4$ and remembering that $\lambda_{24}^{(4)} = 0$, we have $-m\lambda_{34}^{(4)} = -\frac{1}{r}\lambda_{34}^{(4)} + \lambda_{34}^{(1)}$, i.e

$$\lambda_{34}^{(4)} = \frac{1}{r}\lambda_{34}^{(1)}$$

The two relations binding $\lambda_{34}^{(4)}$ and $\lambda_{34}^{(1)}$ now yield $\lambda_{34}^{(4)} = \lambda_{34}^{(1)} = 0$. So w_4 reduces to

$$w_4 = r\lambda_{35}^{(4)}w_{45} + \lambda_{35}^{(4)}w_{35} + \lambda_{56}^{(4)}w_{56} + r\lambda_{36}^{(4)}w_{46} + \lambda_{36}^{(4)}w_{36}$$

At this point, it is of interest to derive the following result.

Result 1. *The irreducible matrix representation of degree 5 of $\mathcal{H}_{F,r^2}(5)$ defined by the matrices Q_i 's is not a constituent of the Lawrence-Krammer representation of degree 10 of the BMW algebra of type A_4 .*

Proof. If $\mathcal{W}$ is a subspace of $\mathcal{V}^{(5)}$ instead, then we simply have

$$w_4 = r\lambda_{35}^{(4)}w_{45} + \lambda_{35}^{(4)}w_{35}$$

Then, by looking at the coefficient of w_{34} in $g_3.w_4 = r w_4$, we get $m\lambda_{45}^{(4)} = 0$, so that $\lambda_{35}^{(4)} = \lambda_{45}^{(4)}$. Then $w_4 = 0$, which is impossible. $\square$

Let's go back to the main proof. By looking at the coefficient of w_{34} in $g_3.w_4 = r w_4$ and using the complete expression for w_4 this time, we get: $m r \lambda_{35}^{(4)} + m r^2 \lambda_{36}^{(4)}$, i.e $\lambda_{36}^{(4)} = -\frac{1}{r}\lambda_{35}^{(4)}$. We see that all the coefficients in w_4 except $\lambda_{56}^{(4)}$ are multiples of $\lambda_{35}^{(4)}$. Moreover, we claim that w_4 may not be a multiple of w_{56} . Indeed, recall the formula

$$g_3 g_4 . w_4 = -\frac{1}{r} g_4 . w_4 - \frac{1}{r^2} w_4$$

and observe that in $-\frac{1}{r} g_4 . w_4 - \frac{1}{r^2} w_4$, there is no term in w_{36} while in $g_3 g_4 . w_4$, there is one. Thus, without loss of generality, we may set $\lambda_{35}^{(4)} = 1$. So,

$$w_4 = r w_{45} + w_{35} + \lambda_{56}^{(4)} w_{56} - w_{46} - \frac{1}{r} w_{36}$$

We then deduce a complete expression for w_1 by using the relation $w_1 = g_2.w_4 + \frac{1}{r}w_4$:

$$w_1 = (1 + r^2)w_{45} + r w_{35} + w_{25} + \left(r + \frac{1}{r}\right)\lambda_{56}^{(4)}w_{56} - \left(r + \frac{1}{r}\right)w_{46} - w_{36} - \frac{1}{r}w_{26}$$

A contradiction now arises when looking at the coefficient of w_{26} in $g_3.w_1 = -\frac{1}{r}w_1 + w_4$. Indeed, this yields $-1 = \frac{1}{r^2}$ and contradicts $(r^2)^2 \neq 1$.

Suppose now that there exists a basis $(w_1, w_2, w_3, w_4, w_5)$ of $\mathcal{W}$ in which the matrices of the left action by the g_i 's, $i = 1, \dots, 4$ are the P_i 's. We read on the matrices P_1 and P_3 that $g_1.w_4 = -\frac{1}{r}w_4$ and $g_3.w_4 = -\frac{1}{r}w_4$. Thus, we have:

$$w_4 = \mu_{13}^{(4)} w_{13} + \mu_{23}^{(4)} w_{23} + \mu_{24}^{(4)} w_{24} + \mu_{14}^{(4)} w_{14},$$

where the coefficients are related by $\mu_{14}^{(4)} = \mu_{23}^{(4)} = -\frac{1}{r}\mu_{13}^{(4)} = -r\mu_{24}^{(4)}$. In particular, all these coefficients are nonzero. As $g_1.w_1 = g_3.w_1$, by looking at the coefficient in w_{23} , we obtain

$$\mu_{13}^{(1)} - m\mu_{23}^{(1)} = \mu_{24}^{(1)}$$

Moreover, by looking at the same coefficient w_{23} in $g_3.w_1 = r w_1 + w_4$, we get

$$\mu_{24}^{(1)} = r\mu_{23}^{(1)} + \mu_{23}^{(4)}$$

Combining both equations yields

$$\mu_{13}^{(1)} - \frac{1}{r}\mu_{23}^{(1)} = \mu_{23}^{(4)}$$

Further, by looking at the coefficient in w_{13} in the same equation, we get $0 = r\mu_{13}^{(1)} + \mu_{13}^{(4)}$, as there is no term in w_{14} in w_1 by $g_2.w_1 = -\frac{1}{r}w_1$. Then we get $\mu_{23}^{(1)} = 0$. Looking at the coefficient of w_{23} in $g_2.w_4 = w_1 + r w_4$ now yields $r\mu_{23}^{(4)} = \frac{1}{l}\mu_{23}^{(4)} + \frac{m}{l}\mu_{13}^{(4)}$, which by using the relations from the beginning reads

$$\left(\frac{m}{l} + \frac{1}{r}(1 - \frac{1}{l})\right)\mu_{13}^{(4)} = 0$$

As the coefficient $\mu_{13}^{(4)}$ is nonzero, this forces $l = r$. It will be useful to derive the following result along the way:

Result 2. Assume $\mathcal{H}_{F,r^2}(5)$ is semisimple. If there exists an irreducible 5-dimensional invariant subspace of $\mathcal{V}^{(5)}$ then $l = r$.

Proof. Indeed, if we assume $\mathcal{W} \subset \mathcal{V}^{(5)}$ instead of $\mathcal{W} \subset \mathcal{V}^{(6)}$ in the computations above, they are unchanged and lead to the same conclusion. $\square$

As seen along the way, w_4 is, up to a multiplication by a scalar that can be set to 1 without loss of generality,

$$w_4 = w_{23} + w_{14} - r w_{13} - \frac{1}{r} w_{24}$$

Then the other spanning vectors of $\mathcal{W}$ must be:

$$\begin{aligned} w_5 &= g_4.w_4 &= r w_{23} + w_{15} - r^2 w_{13} - \frac{1}{r} w_{25} \\ w_1 &= g_2.w_4 - r w_4 &= w_{24} - r w_{12} + w_{13} - \frac{1}{r} w_{34} \\ w_2 &= g_4.w_1 &= w_{25} - r^2 w_{12} + r w_{13} - \frac{1}{r} w_{35} \\ w_3 &= g_3.w_2 - r w_2 &= r w_{14} - r^2 w_{13} + w_{35} - \frac{1}{r} w_{45} \end{aligned}$$

where we replaced l by r . But $\mathcal{W}$ is an invariant subspace of $\mathcal{V}^{(6)}$. In particular, it must be invariant under the action by g_5 . This is not compatible with the spanning set above. We conclude that it is impossible to have

$$\mathcal{W} \downarrow_{\mathcal{H}_{F,r^2}(5)} \simeq S^{(3,2)} \quad \text{or} \quad \mathcal{W} \downarrow_{\mathcal{H}_{F,r^2}(5)} \simeq S^{(2,2,1)}$$

and so $\mathcal{W}$ cannot be isomorphic to $S^{(3,3)}$ or $S^{(2,2,2)}$. Thus, by previous work, the existence of an irreducible 5-dimensional invariant subspace of $\mathcal{V}^{(6)}$ implies that $l \in \{\frac{1}{r^3}, -\frac{1}{r^3}\}$. This completes the proof of the Theorem. $\square$

3.4 The cases $l = r$ and $l = -r^3$

In this section, we show that when $l = r$ the representation $\nu^{(n)}$ is reducible for all $n \geq 4$ and when $l = -r^3$, the representation is reducible for all $n \geq 3$. The latter point is true by Theorem 4 when $n = 3$ and by Theorem 5 when $n = 4$. To do so, we show that some proper invariant subspace $K(n)$ of $\mathcal{V}^{(n)}$, defined in the Proposition below, is nontrivial.


Proposition 2. *For any two nodes i and j with $1 \leq i < j \leq n$, define*

$$c_{ij} = \begin{cases} g_{j-1} \cdots g_{i+1} e_i g_{i+1}^{-1} \cdots g_{j-1}^{-1} & \text{if } j \geq i+2 \\ e_i & \text{if } j = i+1 \end{cases}$$

Then, $K(n) = \cap_{1 \leq i < j \leq n} \text{Ker } \nu^{(n)}(c_{ij})$ is a proper invariant subspace of $\mathcal{V}^{(n)}$. Moreover, any proper invariant subspace of $\mathcal{V}^{(n)}$ must be contained in $K(n)$.

Proof. $K(n)$ is proper, as is visible on the expressions for $\nu^{(n)}(e_i)$. Further, if an x_β is annihilated by all the g_i conjugates of the e_i 's, then $\nu_k(x_\beta)$ is also annihilated by these same elements. Verification of this fact is tedious and can be found in [7], §2. Hence $K(n)$ is invariant. Let $\mathcal{W}$ be a proper invariant subspace of $\mathcal{V}^{(n)}$. By Proposition 1, we have $\nu^{(n)}(c_{i,i+1})(\mathcal{W}) = 0$ for all i with $1 \leq i \leq n-1$. This fact is also true for the other conjugates c_{ij} 's. Hence $\mathcal{W}$ must be contained in $K(n)$ $\square$

This is how an element c_{ij} is represented in the tangle algebra:



It has two horizontal strands: one at the top and one at the bottom, each joining nodes i and j and moreover, when $j \geq i+2$, such horizontal strands over-cross all the vertical strands that they intersect.

To show that $\nu^{(n)}$ is reducible, it will suffice to exhibit a nontrivial element in $K(n)$ when $l = r$ or $l = -r^3$. The following Proposition shows that $K(4)$ is nontrivial and irreducible when $l = r$ and $\mathcal{H}_{F,r^2}(4)$ is semisimple.

Proposition 3. *Assume $\mathcal{H}_{F,r^2}(4)$ is semisimple. There exists an irreducible 2-dimensional invariant subspace of $\mathcal{V}^{(4)}$ if and only if $l = r$. If so it is unique and it is $K(4)$. Moreover, it is spanned over F by the two linearly independent vectors:*

$$v_1 = w_{13} - \frac{1}{r} w_{23} + \frac{1}{r^2} w_{24} - \frac{1}{r} w_{14} \quad (28)$$

$$v_2 = w_{12} - \frac{1}{r} w_{13} - \frac{1}{r} w_{24} + \frac{1}{r^2} w_{34} \quad (29)$$

Proof. When $\mathcal{H}_{F,r^2}(4)$ is semisimple, the following three matrices

$$H_1 = \begin{bmatrix} -\frac{1}{r} & 1 \\ 0 & r \end{bmatrix}, \quad H_2 = \begin{bmatrix} r & 0 \\ 1 & -\frac{1}{r} \end{bmatrix}, \quad H_3 = \begin{bmatrix} -\frac{1}{r} & 1 \\ 0 & r \end{bmatrix}$$

define an irreducible matrix representation of degree 2 of $\mathcal{H}_{F,r^2}(4)$. Suppose $\mathcal{W}$ is an irreducible 2-dimensional invariant subspace of $\mathcal{V}^{(4)}$. Then $\mathcal{W}$ has a basis (v_1, v_2) of vectors such that the matrix of ν_i in this basis is H_i . Since $\nu_1(v_1) = -\frac{1}{r}v_1$ (resp $\nu_3(v_1) = -\frac{1}{r}v_1$), there is no term in w_{34} (resp w_{12}) in v_1 and since $\nu_2(v_2) = -\frac{1}{r}v_2$, there is no term in w_{14} in v_2 . Let the λ_{ij} 's (resp μ_{ij} 's) denote the coefficients of the w_{ij} 's in v_1 (resp v_2). We have:

$$\begin{cases} \lambda_{23} = -\frac{1}{r}\lambda_{13} & \text{and} & \lambda_{24} = -\frac{1}{r}\lambda_{14} & \text{by (14) with } i=1 \text{ and } \gamma = -\frac{1}{r} \\ \lambda_{24} = -\frac{1}{r}\lambda_{23} & \text{and} & \lambda_{14} = -\frac{1}{r}\lambda_{13} & \text{by (15) with } i=3 \text{ and } \gamma = -\frac{1}{r} \\ \mu_{34} = -\frac{1}{r}\mu_{24} & & & \text{by (14) with } i=2 \text{ and } \gamma = -\frac{1}{r} \\ \mu_{13} = -\frac{1}{r}\mu_{12} & & & \text{by (15) with } i=2 \text{ and } \gamma = -\frac{1}{r} \end{cases}$$

Hence, without loss of generality, $v_1 = w_{13} - \frac{1}{r}w_{23} + \frac{1}{r^2}w_{24} - \frac{1}{r}w_{14}$ and v_2 is a multiple of $w_{12} - \frac{1}{r}w_{13} + \mu(w_{24} - \frac{1}{r}w_{34}) + \mu'w_{23}$, where μ and μ' are scalars to determine. The relation $\nu_2(v_1) = r v_1 + v_2$ sets $v_2 = w_{12} - \frac{1}{r}w_{13} + \mu(w_{24} - \frac{1}{r}w_{34}) + \mu'w_{23}$, by just looking at the coefficient in w_{12} . The same relation yields $\mu = -\frac{1}{r}$ by looking at the coefficient in w_{24} . Next, by looking at the coefficient of w_{23} in $\nu_3(v_2) = v_1 + r v_2$, we get $r\mu' - \frac{1}{r} = \mu$. Replacing μ by $-\frac{1}{r}$ now yields $\mu' = 0$. We thus get the expressions in (28) and (29) for v_1 and v_2 respectively. Also, by looking at the coefficient in w_{12} in $\nu_1(v_2) = v_1 + r v_2$, we have $\frac{1}{r} - m = r$, hence $l = r$. Conversely, if $l = r$, it is a direct verification that the vectors v_1 and v_2 given by the formulas (28) and (29) are linearly independent and that they verify the relations:

$$\begin{aligned} \nu_i(v_i) &= -\frac{1}{r}v_i \quad \text{when } i \in \{1, 2\} \quad \& \quad \nu_3(v_1) = -\frac{1}{r}v_1 \\ \nu_1(v_2) &= v_1 + r v_2 = \nu_3(v_2) \quad \& \quad \nu_2(v_1) = r v_1 + v_2 \end{aligned}$$

Thus, their linear span over F is an irreducible 2-dimensional invariant subspace of $\mathcal{V}^{(4)}$. It remains to show that it is in fact $K(4)$. By Proposition 2, $\text{Span}_F(v_1, v_2)$ is contained in $K(4)$. If $K(4)$ is three-dimensional, either it is irreducible and so $l \in \{-r^3, \frac{1}{r}, -\frac{1}{r}\}$ by Theorem 5. This is impossible as $l = r$. Or it is reducible and it must contain a one-dimensional invariant subspace. Then by Theorem 4, it forces $l = \frac{1}{r^3}$, which is again impossible. If $K(4)$ is four-dimensional, then $K(4)$ is not irreducible as its dimension is not 1, 2 or 3. Since we just saw that there exists only one irreducible 2-dimensional invariant subspace of $\mathcal{V}^{(4)}$ when $l = r$, $K(4)$ must then contain a one-dimensional invariant subspace, which is again impossible. For similar reasons, it is also impossible to have $k(4) = 5$, hence the only possibility that is left is to have $k(4) = 2$. We conclude that $K(4) = \text{Span}_F(v_1, v_2)$ and $K(4)$ is thus irreducible. $\square$


The next Proposition shows the reducibility of the representation when $l = r$ and $n \geq 4$, where we still assume that $\mathcal{H}_{F,r^2}(n)$ is semisimple.

Proposition 4. Assume $l = r$ and $\mathcal{H}_{F,r^2}(n)$ is semisimple. Then the vector $v_1 = w_{13} - \frac{1}{r}w_{23} + \frac{1}{r^2}w_{24} - \frac{1}{r}w_{14}$ of Proposition 3 belongs to $K(n)$ for all $n \geq 4$. Thus, $\nu^{(n)}$ is reducible when $l = r$ and $n \geq 4$.


Proof. For $n = 4$, the result is contained in Proposition 3. When $i \geq 5$, we simply have for any $j \geq i + 2$, $\nu_{i+1}^{-1} \dots \nu_{j-1}^{-1}(v_1) = \frac{1}{r^{j-i-1}} v_1$ and since $\nu^{(n)}(e_i)(v_1) = 0$, we see that v_1 is thus annihilated by all the $\nu^{(n)}(c_{ij})$ with $i \geq 5$. Also, since we just saw in Proposition 3 that v_1 is in $K(4)$, v_1 is annihilated by all the $\nu^{(n)}(c_{ij})$'s with $j \leq 4$. Thus, it suffices to check that v_1 is annihilated by $\nu^{(n)}(c_{1j}), \nu^{(n)}(c_{2j}), \nu^{(n)}(c_{3j})$ and $\nu^{(n)}(c_{4j})$ for any $j \geq 5$. We will use the following formulas that give the action of the c_{ij} 's on the basis vectors of the L-K space in some relevant cases here:

$$\left| \begin{array}{ll} \nu^{(n)}(c_{ij})(w_{i,j-k}) = \frac{1}{l^{r^{k-1}}} w_{ij} & (R_k)_{1 \leq k \leq j-i-1} \\ \nu^{(n)}(c_{ij})(w_{i-k,i}) = \frac{1}{r^{(k-1)+(j-i-1)}} w_{ij} & (L_{j-i,k})_{1 \leq k \leq i-1} \\ \nu^{(n)}(c_{ij})(w_{i-t,j-s}) = \left(\frac{1}{r^{t+s-1}} - \frac{1}{r^{t+s-3}} \right) \left(\frac{1}{l} - \frac{1}{r} \right) w_{ij} & (C_{t,s})_{1 \leq t \leq i-1, 1 \leq s \leq j-i-1} \end{array} \right.$$

These formulas can be shown and pictured easily by using the tangles. Let's take an example in $\mathcal{V}^{(12)}$. The product tangle $c_{4,9} w_{2,7}$ as represented in the figure below



expands as follows, where we use the Kauffman skein relation:



After doing a Reidemeister's move of type II (as it is described in §2.2 of [9]) on the first tangle of the sum above, we see that it is zero. After "delooping" the second tangle of the sum and using Reidemeister's move II twice, we see that it is obtained from the basis vector $w_{4,9}$ with non-crossed vertical strands

by multiplying it to the right by $g_4^{-1}g_8^{-1}$. The resulting coefficient is $\frac{m}{lr^2} = \frac{1}{l}(\frac{1}{r^3} - \frac{1}{r})$. If we call over-crossing a multiplication at the bottom by a g_i in H and under-crossing a multiplication at the bottom by a g_i^{-1} in H , we see that in the last tangle of the sum, there are five under-crossings and two over-crossings. Thus, the resulting coefficient is $-\frac{mr^2}{r^5} = \frac{1}{r^2} - \frac{1}{r^4}$. When adding these two coefficients, we get $(\frac{1}{r^3} - \frac{1}{r})(\frac{1}{l} - \frac{1}{r})$, which is the coefficient in $(C_{2,2})$. Using this example as a support, it is easy to see that, more generally, for any fixed nodes i and j with $1 \leq i < j \leq n$, the coefficient of $\nu^{(n)}(c_{ij})(w_{i-t,j-s})$, where $1 \leq t \leq i-1$ and $1 \leq s \leq j-i-1$, is:

$$m \left(\frac{1}{l} \frac{1}{r^{(t-1)+(s-1)}} - \frac{r^{\{(j-s)-i\}-1}}{r^{(t-1)+(j-i-1)}} \right)$$

which after simplification is the coefficient in $(C_{t,s})$. Thus, we obtain the family of equations $(C_{t,s})_{1 \leq t \leq i-1, 1 \leq s \leq j-i-1}$. Similarly for fixed nodes i and j , the two families of equations $(R_k)_{1 \leq k \leq j-i-1}$ and $(L_{j-i,k})_{1 \leq k \leq i-1}$ can be pictured easily by using the tangles, or can be established by using the definition of c_{ij} and the expression for the representation in §2.3.2.

When $l = r$, we note that the action of $c_{i,j}$ on $w_{i-t,j-s}$ is zero. From there, we have for $j \geq 5$, where we replaced l by r :

$$\begin{cases} \nu^{(n)}(c_{1,j})(v_1) = \nu^{(n)}(c_{1,j})(w_{13} - \frac{1}{r} w_{14}) = 0 & \text{by } (R_{j-3}) \text{ and } (R_{j-4}) \\ \nu^{(n)}(c_{2,j})(v_1) = \nu^{(n)}(c_{2,j})(-\frac{1}{r} w_{23} + \frac{1}{r^2} w_{24}) = 0 & \text{by } (R_{j-3}) \text{ and } (R_{j-4}) \\ \nu^{(n)}(c_{3,j})(v_1) = \nu^{(n)}(c_{3,j})(w_{13} - \frac{1}{r} w_{23}) = 0 & \text{by } (L_{j-3,2}) \text{ and } (L_{j-3,1}) \\ \nu^{(n)}(c_{4,j})(v_1) = \nu^{(n)}(c_{4,j})(\frac{1}{r^2} w_{24} - \frac{1}{r} w_{14}) = 0 & \text{by } (L_{j-4,2}) \text{ and } (L_{j-4,3}) \end{cases}$$

So v_1 is in $K(n)$ for all $n \geq 4$, as announced. It will be useful to note on the way that by the game of the coefficients, the equalities to the right of the first two lines of equations still hold when $l = -r^3$. $\square$

When $l = -r^3$, we have a similar result. This is the object of the next proposition.

Proposition 5. *Assume $\mathcal{H}_{F,r^2}(n)$ is semisimple. When $l = -r^3$, the vector u_1 defined as in Theorem 5 by $u_1 = r w_{23} + w_{13} + (\frac{1}{r} + \frac{1}{r^3})w_{34} - w_{24} - \frac{1}{r} w_{14}$ belongs to $K(n)$ for all $n \geq 4$. Thus, when $l = -r^3$, the representation $\nu^{(n)}$ is reducible for every $n \geq 3$.*

Proof. When $l = -r^3$, $\nu^{(3)}$ is reducible by Theorem 4 and $\nu^{(4)}$ is also reducible by Theorem 5. Suppose now $n \geq 5$. To show that u_1 is in $K(n)$, like in the case $l = r$, it will suffice to check that $\nu^{(n)}(c_{ij})(u_1) = 0$ for all $i \leq 4$ and $j \geq 5$. With $l = -r^3$, the coefficients of type $(C_{t,s})$ are no longer zero. But we have: $\nu^{(n)}(c_{2,j})(w_{13} - \frac{1}{r} w_{14}) = 0$ by $(C_{1,j-3})$ and $(C_{1,j-4})$. For $\nu^{(n)}(c_{3,j})(u_1)$, there is no shortcut and a complete evaluation must be performed. We have, where we respected the same order of the terms in Proposition 5 for the coefficients:

$$\begin{aligned} \nu^{(n)}(c_{3,j})(u_1) &= r \frac{1}{r^{j-4}} + \frac{1}{r^{j-3}} + \left(\frac{1}{r} + \frac{1}{r^3} \right) \left(-\frac{1}{r^3 r^{j-5}} \right) \\ &\quad + \left(\frac{1}{r^{j-4}} - \frac{1}{r^{j-6}} \right) \left(\frac{1}{r^3} + \frac{1}{r} \right) + \frac{1}{r} \left(\frac{1}{r^{j-3}} - \frac{1}{r^{j-5}} \right) \left(\frac{1}{r^3} + \frac{1}{r} \right) w_{3,j} \end{aligned}$$

The rules used are, in the same order: $(L_{j-3,1})$, $(L_{j-3,2})$, (R_{j-4}) , $(C_{1,j-4})$ and $(C_{2,j-4})$. All the coefficients cancel nicely to give $\nu^{(n)}(c_{3,j})(u_1) = 0$.

Finally, for $\nu^{(n)}(c_{4,j})(u_1)$, only the terms in u_1 whose last node is node number 4 yield a nonzero contribution, the first one contributing with a coefficient $(\frac{1}{r} + \frac{1}{r^3}) \frac{1}{r^{j-5}}$, the second one with a coefficient $-\frac{1}{r^{j-4}}$ and the third one with a coefficient $-\frac{1}{r} \frac{1}{r^{j-3}}$ by rules $(L_{j-4,1})$, $(L_{j-4,2})$ and $(L_{j-4,3})$ respectively. The sum of these three coefficients is zero.

Thus, we are done with all the cases and u_1 belongs to $K(n)$ for all $n \geq 4$. $\square$
At this stage, we have shown that when l and r take the values of Theorem 1, the representation $\nu^{(n)}$ is reducible. In the next section, we show conversely that if $\nu^{(n)}$ is reducible, then l and r must be related in the way described in Theorem 1.

4 Proof of the main theorem

We recall from Proposition 2 that any proper irreducible invariant subspace of $\mathcal{V}^{(n)}$ is an irreducible $\mathcal{H}_{F,r^2}(n)$ -module. When $n = 3$, the irreducible $\mathcal{H}_{F,r^2}(3)$ -modules have dimension 1 or 2. We showed in Theorem 4 that there exists a one-dimensional invariant subspace of $\mathcal{V}^{(3)}$ if and only if $l \in \{r^3, \frac{1}{r^3}\}$ and we saw in Theorem 5 that there exists an irreducible 2-dimensional invariant subspace of $\mathcal{V}^{(3)}$ if and only if $l \in \{1, -1\}$. Hence the main theorem is proven for $n = 3$. When $n = 4$, the irreducible $\mathcal{H}_{F,r^2}(4)$ -modules have dimensions 1, 2 or 3. By Theorem 4 (resp Theorem 5, resp Proposition 3), there exists a one-dimensional (resp an irreducible 2-dimensional, resp an irreducible 3-dimensional) invariant subspace of $\mathcal{V}^{(4)}$ if and only if $l = \frac{1}{r^5}$ (resp $l = r$, resp $l \in \{-r^3, \frac{1}{r}, -\frac{1}{r}\}$). Thus, the main theorem also holds for $n = 4$. Suppose now $n \geq 5$. By §3, it suffices to prove that if $\nu^{(n)}$ is reducible, then $l \in \{r, -r^3, \frac{1}{r^{2n-3}}, \frac{1}{r^{n-3}}, -\frac{1}{r^{n-3}}\}$ and the proof of the main theorem will be complete. We show it for $n = 5$ and $n = 6$, then proceed by induction on n .

4.1 The case $n = 5$

If $\mathcal{W}$ is an irreducible proper invariant subspace of $\mathcal{V}^{(5)}$, then $\dim(\mathcal{W}) \in \{1, 4, 5, 6\}$, as $\mathcal{W}$ is an irreducible $\mathcal{H}_{F,r^2}(5)$ -module by Corollary 1 of §3.1. If $\dim(\mathcal{W}) = 5$, it forces $l = r$ by Result 2. If $\dim(\mathcal{W}) = 1$, it forces $l = \frac{1}{r^7}$ by Theorem 4 and if $\dim(\mathcal{W}) = 4$, it forces $l \in \{\frac{1}{r^2}, -\frac{1}{r^2}\}$ by Theorem 5. From now on, assume that $l \notin \{r, \frac{1}{r^7}, \frac{1}{r^2}, -\frac{1}{r^2}\}$. We will show that $l = -r^3$. By choice for l and r , we have $\dim(\mathcal{W}) = 6$. Then $\mathcal{W} \cap \mathcal{V}^{(4)} \neq \{0\}$, as otherwise $\mathcal{W} \oplus \mathcal{V}^{(4)} \subseteq \mathcal{V}^{(5)}$, which implies on the dimensions $\dim \mathcal{W} \leq 10 - 6 = 4$. Since $\mathcal{W} \cap \mathcal{V}^{(4)}$ is a proper invariant subspace of $\mathcal{V}^{(4)}$, the representation $\nu^{(4)}$ is then reducible, which yields $l \in \{-r^3, \frac{1}{r}, -\frac{1}{r}, \frac{1}{r^5}\}$. We will show that it is impossible to have $l \in \{\frac{1}{r}, -\frac{1}{r}, \frac{1}{r^5}\}$, unless $l = \frac{1}{r^5} = -r^3$.

Let's first assume that $l = \frac{1}{r^5}$ and show that under our assumptions, it forces $l = -r^3$. When $l = \frac{1}{r^5}$, there exists a one-dimensional invariant subspace, say $\mathcal{U}_1$, of $\mathcal{V}^{(4)}$ by Theorem 4 and by Proposition 2 it is contained in $K(4)$. In particular the dimension $k(4)$ of $K(4)$ is 1, 2, 3, 4 or 5. But since $\mathcal{W} \cap \mathcal{V}^{(4)}$ is a proper invariant subspace of $\mathcal{V}^{(4)}$, it must be contained in $K(4)$ by Proposition 2. Hence we have $k(4) \geq \dim(\mathcal{W} \cap \mathcal{V}^{(4)})$. Also, since $\dim(\mathcal{W} \cap \mathcal{V}^{(4)}) = \dim(\mathcal{W}) + \dim(\mathcal{V}^{(4)}) - \dim(\mathcal{W} + \mathcal{V}^{(4)}) \geq 12 - \dim(\mathcal{V}^{(5)}) = 2$, we get $k(4) \geq 2$. By semisimplicity of $\mathcal{H}_{F,r^2}(4)$ and by uniqueness of a one-dimensional invari-

ant subspace of $\mathcal{V}^{(4)}$ when it exists, $\mathcal{U}_1$ then has a summand S in $K(4)$ that is of dimension greater than or equal to 2. S is an invariant subspace of $\mathcal{V}^{(4)}$ and it cannot contain a one-dimensional invariant subspace. Nor can it contain an irreducible 2-dimensional invariant subspace by Proposition 3 since we assumed $l \neq r$. Also, since $1 + \dim(S) = k(4) \leq 5$, we note that $\dim(S) \leq 4$. Then, by the same arguments already exposed, $\dim(S) = 3$ and S is irreducible. By Theorem 5, we now get $l \in \{-\frac{1}{r}, \frac{1}{r}, -r^3\}$. But since $l = \frac{1}{r^5}$, it forces $l = -r^3$ as it is impossible to have $(r^2)^4 = 1$ when $\mathcal{H}_{F,r^2}(5)$ is semisimple.

Assume next that $l \in \{\frac{1}{r}, -\frac{1}{r}\}$. We show that these values lead to a contradiction. First, by choice for l and r and Theorem 5 (case $n = 4$), $\mathcal{V}^{(4)}$ contains an irreducible 3-dimensional invariant subspace and by Proposition 2, this proper invariant subspace must be contained in $K(4)$. Hence $k(4) \geq 3$. Since there cannot exist any one-dimensional invariant subspace of $\mathcal{V}^{(4)}$ (as $l \neq \frac{1}{r^5}$ when $(r^2)^4 \neq 1$) or any irreducible 2-dimensional invariant subspace of $\mathcal{V}^{(4)}$ (as $l \neq r$), we cannot have $k(4) \in \{4, 5\}$. Thus, we have $k(4) = 3$ and so $K(4)$ is irreducible. Now the irreducibility of $K(4)$ and the fact that $0 \subset \mathcal{W} \cap \mathcal{V}^{(4)} \subseteq K(4)$ implies that

$$\mathcal{W} \cap \mathcal{V}^{(4)} = K(4) \quad (30)$$

We show that this is impossible.

Consider first the case when $l = \frac{1}{r}$. When $l = \frac{1}{r}$, the vector $w_{14} - w_{23}$ belongs to $K(4)$. Indeed, this vector is

$$\frac{1}{r^2 + \frac{1}{r^2}} \left(r v_1^{(4)} + \left(r - \frac{1}{r}\right) v_2^{(4)} - \frac{1}{r} v_3^{(4)} \right)$$

Then it also belongs to $\mathcal{W}$. It follows that $\nu^{(5)}(e_4)(w_{14} - w_{23}) = \frac{1}{r^2} x_{\alpha_4} \in \mathcal{W}$, as $\mathcal{W}$ is an invariant subspace of $\mathcal{V}^{(5)}$. But then, by the same argument as in the proof of Proposition 1, $\mathcal{W}$ is the whole space $\mathcal{V}^{(5)}$, in contradiction with $\mathcal{W}$ is proper. Thus, it is impossible to have (30) and so l cannot take the value $\frac{1}{r}$.

Consider now the case when $l = -\frac{1}{r}$. The vector $w_{14} - w_{23} + w_{12} + w_{34}$ belongs to $K(4)$ since it is

$$\frac{1}{r + \frac{1}{r}} \left(v_1^{(4)} + v_3^{(4)} \right)$$

By (30), this vector also belongs to $\mathcal{W}$. But then

$$\nu^{(5)}(e_4)(w_{14} - w_{23} + w_{12} + w_{34}) = \left(1 + \frac{1}{r^2}\right) x_{\alpha_4} \in \mathcal{W}$$

As $(r^2)^2 \neq 1$, this implies in turn that x_{α_4} is in $\mathcal{W}$. Then $\mathcal{W}$ is the whole space $\mathcal{V}^{(5)}$, a contradiction. Thus, it is also impossible to have $l = -\frac{1}{r}$.

We have now shown that if $\nu^{(5)}$ is reducible and $l \notin \{r, \frac{1}{r}, \frac{1}{r^2}, -\frac{1}{r^2}\}$, then $l = -r^3$. Thus, if $\nu^{(5)}$ is reducible, then $l \in \{r, -r^3, \frac{1}{r}, -\frac{1}{r^2}, \frac{1}{r^2}\}$.

4.2 The case $n = 6$

Let $\mathcal{W}$ be an irreducible proper invariant subspace of $\mathcal{V}^{(6)}$. So $\mathcal{W}$ is an irreducible $\mathcal{H}_{F,r^2}(6)$ -module. The irreducible representations of $\mathcal{H}_{F,r^2}(6)$ have degrees 1, 5, 9, 10, 16. The vector space $\mathcal{V}^{(6)}$ is 15-dimensional. Hence $\dim(\mathcal{W}) \in$

$\{1, 5, 9, 10\}$. If $\dim(\mathcal{W}) = 1$, then $l = \frac{1}{r^9}$ by Theorem 4 and if $\dim(\mathcal{W}) = 5$, then $l \in \{\frac{1}{r^3}, -\frac{1}{r^3}\}$ by Theorem 5. Suppose now $l \notin \{\frac{1}{r^9}, \frac{1}{r^3}, -\frac{1}{r^3}\}$. Then $\dim(\mathcal{W}) \geq 9$, which implies in particular that $\mathcal{W} \cap \mathcal{V}^{(5)} \neq \{0\}$. Moreover, $\mathcal{W} \cap \mathcal{V}^{(5)}$ is a proper subspace of $\mathcal{V}^{(5)}$, as otherwise $\mathcal{W}$ would contain $\mathcal{V}^{(5)}$ and would in fact be the whole space $\mathcal{V}^{(6)}$. Hence we see that $\nu^{(5)}$ is reducible, which implies that

$$l \in \left\{ r, -r^3, \frac{1}{r^7}, \frac{1}{r^2}, -\frac{1}{r^2} \right\} \quad (31)$$

by the case $n = 5$. Also, if $\mathcal{W} \cap \mathcal{V}^{(4)} = \{0\}$, then $\mathcal{W} \oplus \mathcal{V}^{(4)} \subseteq \mathcal{V}^{(6)}$ and so $\dim(\mathcal{W}) \leq 15 - 6 = 9$. Then $\dim \mathcal{W} = 9$. We notice that $\dim(\mathcal{W}) + \dim(\mathcal{V}^{(4)}) = \dim(\mathcal{V}^{(6)})$. Thus, we get $\mathcal{W} \oplus \mathcal{V}^{(4)} = \mathcal{V}^{(6)}$. But since $\mathcal{W} \subseteq K(6)$ by Proposition 2, we must have in particular $\nu^{(6)}(e_5)(\mathcal{W}) = 0$. But e_5 also acts trivially on $\mathcal{V}^{(4)}$. It follows that e_5 acts trivially on the direct sum $\mathcal{W} \oplus \mathcal{V}^{(4)}$, hence acts trivially on $\mathcal{V}^{(6)}$. This is a contradiction. Hence, we have $\mathcal{W} \cap \mathcal{V}^{(4)} \neq \{0\}$. Also, $\mathcal{W} \cap \mathcal{V}^{(4)}$ is a proper invariant subspace of $\mathcal{V}^{(4)}$. Consequently, $\nu^{(4)}$ is reducible and by the case $n = 4$, we have

$$l \in \left\{ r, -r^3, \frac{1}{r^5}, \frac{1}{r}, -\frac{1}{r} \right\} \quad (32)$$

Since $r^2 \neq 1$, $(r^2)^3 \neq 1$ and $(r^2)^6 \neq 1$ when $\mathcal{H}_{F,r^2}(6)$ is semisimple, (31) and (32) imply that $l \in \{r, -r^3\}$. Thus, if $\nu^{(6)}$ is reducible and $l \notin \{\frac{1}{r^9}, \frac{1}{r^3}, -\frac{1}{r^3}\}$, then $l \in \{r, -r^3\}$. So if $\nu^{(6)}$ is reducible, then $l \in \{r, -r^3, \frac{1}{r^9}, \frac{1}{r^3}, -\frac{1}{r^3}\}$.

4.3 Proof of the main theorem when $n \geq 7$

By the work from previous parts, the main theorem holds for $n \in \{3, 4, 5, 6\}$. When $n \geq 7$, we proceed by induction to prove the theorem. Given an integer n with $n \geq 7$, suppose the main theorem holds for $\nu^{(n-1)}$ and $\nu^{(n-2)}$. We already saw that when $l \in \{r, -r^3, \frac{1}{r^{2n-3}}, \frac{1}{r^{n-3}}, -\frac{1}{r^{n-3}}\}$, the representation $\nu^{(n)}$ is reducible. We will show conversely that if $\nu^{(n)}$ is reducible, it forces these values for l and r . The theorem will then be proven. Suppose $\nu^{(n)}$ is reducible and let $\mathcal{W}$ be an irreducible nontrivial proper invariant subspace of $\mathcal{V}^{(n)}$. By Corollary 1, we know that $\mathcal{W}$ is an irreducible $\mathcal{H}_{F,r^2}(n)$ -module. The following proposition is part of the author's work in [4].

Proposition 6. *Let K be a field of characteristic zero. Let n be an integer with $n \geq 9$. Every irreducible $K \operatorname{Sym}(n)$ -module is either isomorphic to one of the Specht modules $S^{(n)}, S^{(n-1,1)}, S^{(n-2,2)}, S^{(n-2,1,1)}$ or to one of their conjugates, or has dimension greater than $\frac{(n-1)(n-2)}{2}$.*

We have the Corollary on the dimensions:

Corollary 2. *Assume $\mathcal{H}_{F,r^2}(n)$ is semisimple.*

(i) *Let $\mathcal{D}$ be an irreducible $F \operatorname{Sym}(n)$ -module with $n = 7$ or $n \geq 9$, where F is a field of characteristic zero. Then, there are two possibilities:*

$$\text{either } \dim \mathcal{D} \in \left\{ 1, n-1, \frac{n(n-3)}{2}, \frac{(n-1)(n-2)}{2} \right\}$$

$$\text{or } \dim \mathcal{D} > \frac{(n-1)(n-2)}{2}$$

- (ii) Let $\mathcal{D}$ be an irreducible $F \operatorname{Sym}(8)$ -module, where F is a field of characteristic zero. Then $\dim \mathcal{D} \in \{1, 7, 14, 20, 21\}$ or $\dim \mathcal{D} > 21$.
- (iii) The first two points hold if $\mathcal{D}$ is an irreducible $\mathcal{H}_{F,r^2}(n)$ -module.

Proof. (ii) can be seen directly by using the Hook formula. Point (i) is for $n \geq 9$ a direct consequence of Proposition 6 after noticing that $S^{(n-2,2)}$ has dimension $\frac{n(n-3)}{2}$ and $S^{(n-2,1,1)}$ dimension $\frac{(n-1)(n-2)}{2}$. When $n = 7$, the statement also holds by direct investigation, using for instance the Hook formula. In characteristic zero, when the Hecke algebra $\mathcal{H}_{F,r^2}(n)$ is semisimple, the dimensions of the irreducible $F \operatorname{Sym}(n)$ -modules are the same as the dimensions of the irreducible $\mathcal{H}_{F,r^2}(n)$ -modules, hence (iii). $\square$

Let's go back to the proof of the Main Theorem. Suppose first $n = 7$ or $n \geq 9$. So $\dim \mathcal{W} \in \{1, n-1, \frac{n(n-3)}{2}, \frac{(n-1)(n-2)}{2}\}$ or $\dim \mathcal{W} > \frac{(n-1)(n-2)}{2}$. First, if $\dim \mathcal{W} = 1$, Theorem 4 implies that $l = \frac{1}{r^{2n-3}}$. Also, if $\dim \mathcal{W} = n-1$, Theorem 5 implies that $l \in \{\frac{1}{r^{n-1}}, -\frac{1}{r^{n-1}}\}$. Suppose $l \notin \{\frac{1}{r^{2n-3}}, \frac{1}{r^{n-3}}, -\frac{1}{r^{n-3}}\}$. Then we have $\dim \mathcal{W} \geq \frac{n(n-3)}{2}$.

Claim 2. Let $\mathcal{W}$ be a subspace of $\mathcal{V}^{(n)}$.

If $\dim \mathcal{W} > n-1$, then $\mathcal{W} \cap \mathcal{V}^{(n-1)} \neq \{0\}$.

If $\dim \mathcal{W} > 2n-3$, then $\mathcal{W} \cap \mathcal{V}^{(n-2)} \neq \{0\}$

Proof. If $\mathcal{W} \cap \mathcal{V}^{(n-1)} = \{0\}$, the L-K space $\mathcal{V}^{(n)}$ contains the direct sum $\mathcal{W} \oplus \mathcal{V}^{(n-1)}$, which yields on the dimensions: $\dim \mathcal{W} + \frac{(n-1)(n-2)}{2} \leq \frac{n(n-1)}{2}$. Then $\dim \mathcal{W} \leq n-1$. Similarly, if $\mathcal{W} \cap \mathcal{V}^{(n-2)} = \{0\}$, we get $\dim \mathcal{W} \leq \frac{n(n-1)}{2} - \frac{(n-2)(n-3)}{2} = 2n-3$. $\square$

Lemma 4. When $n > 6$, we have $\frac{n(n-3)}{2} > 2n-3$ and $\frac{n(n-3)}{2} > n-1$.

By the claim and the lemma, the intersections $\mathcal{W} \cap \mathcal{V}^{(n-1)}$ and $\mathcal{W} \cap \mathcal{V}^{(n-2)}$ are nontrivial. Since $\mathcal{W}$ is proper in $\mathcal{V}^{(n)}$, $\mathcal{W}$ cannot contain $\mathcal{V}^{(n-1)}$. Nor can it contain $\mathcal{V}^{(n-2)}$. Hence $\mathcal{W} \cap \mathcal{V}^{(n-1)}$ (resp $\mathcal{W} \cap \mathcal{V}^{(n-2)}$) is a proper nontrivial invariant subspace of $\mathcal{V}^{(n-1)}$ (resp $\mathcal{V}^{(n-2)}$). Now $\nu^{(n-1)}$ and $\nu^{(n-2)}$ are both reducible. Since we assumed the main theorem to be true for $\nu^{(n-1)}$ and $\nu^{(n-2)}$, we get:

$$l \in \left\{ r, -r^3, \frac{1}{r^{2n-5}}, \frac{1}{r^{n-4}}, -\frac{1}{r^{n-4}} \right\} \cap \left\{ r, -r^3, \frac{1}{r^{2n-7}}, \frac{1}{r^{n-5}}, -\frac{1}{r^{n-5}} \right\}$$

Since $r^2 \neq 1$, $r^{2(n-3)} \neq 1$ and $r^{2n} \neq 1$ when $\mathcal{H}_{F,r^2}(n)$ is semisimple, it only leaves the possibility $l \in \{r, -r^3\}$.

When $n = 8$, if $l \notin \{\frac{1}{r^{13}}, \frac{1}{r^5}, -\frac{1}{r^5}\}$, then we have $\dim \mathcal{W} \geq 14 > 13 = 2 \times 8 - 3$. Hence the same method applies and yields again $l \in \{r, -r^3\}$.

Thus, we have shown that if the representation is reducible and $l \notin \{\frac{1}{r^{2n-3}}, \frac{1}{r^{n-3}}, -\frac{1}{r^{n-3}}\}$, then $l \in \{r, -r^3\}$. $\square$

5 Non semisimplicity of the BMW algebra for some specializations of its parameters

In § 2.1, we let the Hecke algebra H act on the base field F by $g_i.1 = r$ for all $i \in \{3, \dots, n-1\}$. If we consider the action given by $g_i.1 = -\frac{1}{r}$ instead, F is

again a left H -module for this action and we get another left B -module of dimension $\frac{n(n-1)}{2}$ by considering again the tensor product $B_1 \otimes_H F$. We call this representation the conjugate L-K representation. By the symmetry of the roles played by r and $-\frac{1}{r}$, when $n \geq 4$ and $\mathcal{H}_{F,r^2}(n)$ is semisimple, the conjugate L-K representation is reducible exactly when $l \in \{-\frac{1}{r}, \frac{1}{r^3}, -r^{2n-3}, r^{n-3}, -r^{n-3}\}$. In particular, when $n \geq 6$, since $\frac{1}{r^3} \notin \{r, -r^3, \frac{1}{r^{2n-3}}, \frac{1}{r^{n-3}}, -\frac{1}{r^{n-3}}\}$, the two representations are not equivalent. This is also true when $n \in \{4, 5\}$. For instance, for the L-K representation, the trace of the matrix of the left action by g_{n-1} is $\frac{(n-2)(n-3)}{2} r + \frac{1}{l} - (n-2)m$. For the conjugate representation it is $\frac{(n-2)(n-3)}{2} (-\frac{1}{r}) + \frac{1}{l} - (n-2)m$.

We note that Proposition 1 remains valid for the conjugate L-K representation. A consequence of this Proposition is that when the representation is reducible, it is indecomposable. Then the BMW algebra is not semisimple for the values of l and r for which the L-K representation or its conjugate representation are reducible. This is the statement of Theorem 2.

6 Conclusion and future developments

In [2], it is established that I_1/I_2 is generically semisimple where I_1 is the two-sided ideal Be_1B and I_2 the two-sided ideal generated by all the products $e_i e_j$ with $|i - j| > 2$. For each irreducible representation of the Hecke algebra of type A_{n-2} of degree θ , the authors build a generically irreducible representation of B/I_2 of degree $\frac{n(n-1)}{2}\theta$ and show that these are all the inequivalent generically irreducible representations of I_1/I_2 . One of the two so-built inequivalent representations of B/I_2 of degree $\frac{n(n-1)}{2}$ is the Lawrence-Krammer representation. The other one is obtained from the first one by replacing r by $-\frac{1}{r}$. Since the representation $B_1 \otimes_H F$ built in this paper is a generically irreducible representation of B/I_2 and its kernel does not contain I_1 , it must be equivalent to the Lawrence-Krammer representation. We think that the other generically irreducible I_1/I_2 -modules are these with F replaced by an irreducible H -module and that by studying these representations we could show that I_1/I_2 is semisimple if and only if l and r don't take the specializations of the Main Theorem.

References

- [1] N. Bourbaki, Groupes et algebres de Lie, Chapitres 4, 5 et 6, Masson 1981.
- [2] A.M. Cohen, D.A.H. Gijsbers and D.B. Wales, BMW algebras of simply laced type, J. Algebra, 285 (2005) no.2 439 – 450
- [3] A.M. Cohen and D.B. Wales, Linearity of Artin groups of finite type, Isr. J. Math. 131 (2002) 101 – 123
- [4] G.D. James, On the minimal dimensions of irreducible representations of symmetric groups, Math. Proc. Camb. Phil. Soc. 94 (1983) 417 – 424

- [5] D. Krammer, Braid groups are linear, *Annals of Math.*, 155 (2002) 131 – 156
- [6] R. Lawrence, Homological representations of the Hecke algebra, *Comm. Math. Physics*, 135 (1990) 141 – 191.
- [7] C. Levaillant, Irreducibility of the Lawrence-Krammer representation of the BMW algebra of type A_{n-1} , PhD dissertation 2008, California Institute of Technology, <http://etd.caltech.edu/etd/available/etd-05292008-110016/unrestricted/thesis.pdf>
- [8] A. Mathas, Iwahori-Hecke algebras and Schur algebras of the symmetric group, *University Lecture Series of the AMS*, Vol 15
- [9] H.R. Morton and A.J. Wassermann, A basis for the Birman-Wenzl algebra, Preprint 1989
- [10] H. Rui and M. Si, Gram determinants and semisimple criteria for Birman-Wenzl algebras, *ArXiv:math/0607266*, to appear in *J. Reine. Angew. Math.*
- [11] H. Wenzl, Quantum groups and subfactors of type B , C , and D , *Commun. Math. Phys.* 133 (1990) 383 – 432